

# Logical Observation Identifiers Names and Codes (LOINC<sup>®</sup>) Users' Guide

Edited by:

Clem McDonald, MD, Stan Huff, MD, Kathy Mercer, Jo Anna Hernandez,  
Daniel J. Vreeman, PT, DPT

Please send questions and comments to:

LOINC  
c/o Regenstrief Institute, Inc  
410 West 10th St. Suite 2000  
Indianapolis, IN 46202

or via email:

[loinc@regenstrief.org](mailto:loinc@regenstrief.org)

This and other relevant documents and files are available at

<http://loinc.org>

## List of Files:

<b>Description</b>	<b>Format</b>	<b>File Name</b>
LOINC table (database)	MDB	LOINCDB.MDB
LOINC table (database)	ASCII	LOINCDB.TXT
LOINC Users' Guide	PDF	LOINCManual.pdf
RELMA Program		Setup.exe
RELMA Users' Manual	PDF	RELMAManual.pdf

## Table of Contents

<b>1</b>	<b>Introduction .....</b>	<b>4</b>
1.1	Successes .....	4
1.2	What is not part of the name .....	5
1.3	Scope of LOINC .....	6
1.4	The LOINC Code Identifier .....	7
<b>2</b>	<b>Major “Parts” of a Test/Observation Name .....</b>	<b>7</b>
2.1	General naming conventions .....	8
2.1.1	Abbreviations in names of component/analyte .....	8
2.1.2	General naming rules for the component (analyte) part of the fully specified name .....	9
2.1.3	Punctuation in analyte names .....	10
2.1.4	Case insensitivity .....	10
2.1.5	Roman numerals vs. Arabic numerals .....	11
2.2	Component/analyte (1st part) .....	11
2.2.1	Analyte Name (1st subpart) .....	11
2.2.2	Challenge test (2nd subpart) .....	12
2.2.3	Adjustments/corrections (3rd subpart) .....	17
2.2.4	Distinguishing multiple values for any test via the test name (4th subpart) .....	17
2.3	Kind of Property (also called kind of quantity) (2nd part) .....	18
2.4	Time Aspect (Point or moment in time vs. time interval) (3rd part) .....	21
2.4.1	Time Aspect Modifier .....	23
2.5	System (Sample) Type (4th part) .....	24
2.5.1	Super system (2nd subpart) .....	26
2.6	Type of Scale (5th part) .....	26
2.7	Type of Method (6th part) .....	28
2.7.1	DNA/RNA probes/measures .....	29
2.7.2	Immunofluorescence (IF) .....	29
2.7.3	Immune Stain .....	30
2.7.4	Enzyme Immunoassay (EIA) .....	30
2.7.5	Coagulation .....	30
2.7.6	Stains .....	30
2.7.7	Clinical measures .....	30
2.7.8	Imaging studies .....	30
2.8	Short Convenient Names .....	30
2.9	Long Common Names .....	31
2.10	LOINC term names in HL7 messages .....	31
2.11	Classes .....	32
<b>3</b>	<b>Special Cases .....</b>	<b>32</b>
3.1	Findings viewed as variables or as values .....	32
3.1.1	Value .....	32
3.1.2	Variable (Multiple Choice) Approach .....	32
3.2	Blood bank .....	33
3.2.1	Panel reporting: .....	33
3.2.2	Multiple answer reporting: .....	34
3.3	Immunocompetence studies (flow cytometry) .....	34

3.4	General approach to microbiology results .....	35
3.5	Antimicrobial susceptibilities .....	37
3.6	Cell counts .....	37
3.7	Skin tests.....	38
3.8	Toxicology – Drug of Abuse Screening and Confirmation .....	38
3.8.1	Toxicology drug groups .....	39
3.8.2	Cutoffs .....	40
3.8.3	Reporting the method used for screen and confirm.....	41
3.8.4	Individual drug/metabolite test results .....	41
3.8.5	Naming issues.....	42
3.8.6	Summary.....	42
3.9	Molecular Genetics LOINC Naming.....	42
3.9.1	Introduction.....	42
3.9.2	Terminology.....	43
3.9.3	General Molecular genetics naming rules .....	44
3.9.4	Infectious Diseases.....	45
3.9.5	Genetic Diseases .....	45
3.9.6	Hematopathology gene re-arrangement .....	48
3.9.7	Translocations.....	48
3.9.8	Identity testing .....	49
3.9.9	Tumor Relation Tumor Genetics.....	50
3.10	Allergy Testing.....	50
<b>4</b>	<b>Clinical observations and measures .....</b>	<b>51</b>
4.1	Introduction .....	51
4.2	Atomic versus molecular (pre-coordinated names).....	53
4.3	Radiology Reports.....	54
4.3.1	Diagnostic Radiology Reports .....	55
4.3.2	Interventional Radiology Reports .....	61
<b>5</b>	<b>Tumor registry.....</b>	<b>62</b>
<b>6</b>	<b>Claims attachments .....</b>	<b>62</b>
<b>7</b>	<b>HL7 LOINC Document Type Vocabulary Domain.....</b>	<b>62</b>
7.1	Use of document type codes in HL7 messages.....	62
7.2	Relationship with terminologies.....	63
7.3	Elements of Document Type codes .....	64
7.4	Rules for Creating Clinical Notes from Multiple Components .....	70
7.5	Future Work .....	71
<b>8</b>	<b>Order Panels (Batteries) .....</b>	<b>71</b>
8.1	Goals.....	72
8.2	Reflex tests .....	73
8.3	Calculated or derived results .....	73
8.4	Associated observations .....	73
8.5	LOINC Rules for representing order panel names .....	73

9	Evolving principles for naming collections .....	75
9.1	Goals and general approach.....	75
9.2	Collections as orders and observations.....	75
9.3	LOINC SCALE for collections .....	76
9.4	Examples of proposed changes according to new policy .....	76
10	Standardized Assessment Measures .....	76
10.1	Introduction .....	77
10.2	Consolidated Health Informatics endorsement.....	77
10.3	LOINC Representation .....	77
10.3.1	Naming rules and conventions for survey instrument items in LOINC .....	77
10.3.2	Structured answer lists .....	78
<b>11</b>	<b>Editorial Policies and Procedures .....</b>	<b>78</b>
11.1	Concept orientation and LOINC name changes .....	78
11.2	Classification of LOINC term status .....	79
11.3	Concept persistence and term deprecation.....	80
	<b>Appendix A - LOINC Database Structure .....</b>	<b>81</b>
	<b>Appendix B - Classes .....</b>	<b>84</b>
	<b>Appendix C - Calculating Mod 10 Check Digits .....</b>	<b>90</b>
	<b>Appendix D - Procedure for Submitting Additions/Changes to the Database .....</b>	<b>91</b>
	<b>Appendix E - Examples for LOINC Property Matching .....</b>	<b>97</b>
	<b>Appendix F - Acronyms used in LOINC .....</b>	<b>101</b>
	<b>Appendix G - LOINC Committee Members .....</b>	<b>104</b>

**Tables**

<b>Table 1: Hierarchical Structure of Fully Specified Analyte Names .....</b>	<b>7</b>
<b>Table 2: Example Component Abbreviations .....</b>	<b>8</b>
<b>Table 3: Example Case Specifying Conventions .....</b>	<b>11</b>
<b>Table 4: Example Time Delay Post Challenge .....</b>	<b>12</b>
<b>Table 5: Example Challenge Subparts.....</b>	<b>13</b>
<b>Table 6: Example Route Abbreviations for Challenge Part .....</b>	<b>14</b>
<b>Table 7: Example Nature of Challenge.....</b>	<b>15</b>
<b>Table 8: Example LOINC properties.....</b>	<b>19</b>
<b>Table 9: Example Duration Categories .....</b>	<b>23</b>
<b>Table 10: Time Aspect Modifier Codes .....</b>	<b>23</b>
<b>Table 11: Example Laboratory System/Sample Types .....</b>	<b>25</b>
<b>Table 12: Type of Scale .....</b>	<b>26</b>
<b>Table 13: Examples of Method Abbreviations.....</b>	<b>28</b>
<b>Table 14A: Examples of specific methods that would be classed as target amplified DNA/RNA .....</b>	<b>29</b>
<b>Table 14B: Examples of specific methods that would be defined in LOINC as signal amplification methods .....</b>	<b>29</b>

<b>Table 15: Example Culture Results .....</b>	<b>37</b>
<b>Table 16: Drug Susceptibility Methods .....</b>	<b>37</b>
<b>Table 17: Drug of Abuse Methods.....</b>	<b>38</b>
<b>Table 18: Three types of nomenclatures for identifying the location of a genetic defect .....</b>	<b>44</b>
<b>Table 19: List of single letter amino acid codes .....</b>	<b>44</b>
<b>Table 20: Subjects covered to date in clinical LOINC .....</b>	<b>52</b>
<b>Table 21: Examples of Pre-Coordinated Names .....</b>	<b>54</b>
<b>Table 22: Example Clinical Notes .....</b>	<b>63</b>
<b>Table 23. Document Ontology LOINC Naming Rules .....</b>	<b>70</b>
<b>Table 24. Example Document Ontology LOINC Codes.....</b>	<b>71</b>
<b>Table 25: Example Order Sets.....</b>	<b>71</b>
<b>Table 26: Examples of LOINC Panel Names (Order Set Names) .....</b>	<b>74</b>
<b>Table 27: Example of Proposed Changes .....</b>	<b>76</b>
<b>Table 28: LOINC Database Structure .....</b>	<b>81</b>
<b>Table 29: Classes.....</b>	<b>84</b>
<b>Table 30: Example submission .....</b>	<b>92</b>
<b>Table 31a: Access Field Names for Submission .....</b>	<b>93</b>
<b>Table 31b: Content Added by Regenstrief (Fields left blank in submission) .....</b>	<b>Error! Bookmark not defined.</b>
<b>Table 32: Acronyms used in LOINC.....</b>	<b>101</b>

## Figures

<b>Figure 1. Submission Created with Microsoft Access 97 .....</b>	<b>93</b>
<b>Figure 2. An Example Excel Submission (first 9 fields only) .....</b>	<b>94</b>
<b>Figure 3. Example Submission Using ASCII Tab-Delimited File.....</b>	<b>95</b>
<b>Figure 4. The Propose LOINC Form .....</b>	<b>Error! Bookmark not defined.</b>
<b>Figure 5. The Local Code Section.....</b>	<b>Error! Bookmark not defined.</b>
<b>Figure 6. The Similar LOINC Section .....</b>	<b>Error! Bookmark not defined.</b>
<b>Figure 7. Propose LOINC form showing Analyte textbox “dropped down”.....</b>	<b>Error! Bookmark not defined.</b>
<b>Figure 8. Propose LOINC form showing System textbox “dropped up” .....</b>	<b>Error! Bookmark not defined.</b>
<b>Figure 9. Providing comments.....</b>	<b>Error! Bookmark not defined.</b>
<b>Figure 10. Example answers (sample results) .....</b>	<b>Error! Bookmark not defined.</b>
<b>Figure 11. The Answer List Dialog.....</b>	<b>Error! Bookmark not defined.</b>
<b>Figure 12. Defining a new answer list .....</b>	<b>Error! Bookmark not defined.</b>
<b>Figure 13. Highlighting an existing LOINC before proposing a new one.....</b>	<b>Error! Bookmark not defined.</b>
<b>Figure 14. Proposing a LOINC based on an existing one.....</b>	<b>Error! Bookmark not defined.</b>
<b>Figure 15. Review Proposed LOINC Form.....</b>	<b>Error! Bookmark not defined.</b>
<b>Figure 16. Selecting terms the user desires to submit.....</b>	<b>Error! Bookmark not defined.</b>
<b>Figure 17. Windows Common Dialog box used to create LOINC submission files .....</b>	<b>Error! Bookmark not defined.</b>
<b>Figure 18. Message displayed after submission file has been created .....</b>	<b>Error! Bookmark not defined.</b>

## Copyright Notice and License

The LOINC® codes, LOINC® table (regardless of format), LOINC® Release Notes, LOINC® Changes File, and LOINC® Users' Guide are copyright © 1995-2010, Regenstrief Institute, Inc. and the Logical Observation Identifiers Names and Codes (LOINC) Committee. All rights reserved.

The RELMA® program, RELMA® database and associated search index files (subject to the copyright above with respect to the LOINC® codes and LOINC® table included therein), RELMA® Release Notes, and RELMA® Users' Manual are copyright © 1995-2010, Regenstrief Institute, Inc. All rights reserved.

The LOINC® panels and forms file and the LOINC® hierarchies file (subject to the copyright above with respect to the LOINC® codes and LOINC® table to the extent included in the LOINC® panels and forms file and the LOINC® hierarchies file), are copyright © 1995-2010, Regenstrief Institute, Inc. All rights reserved.

LOINC® and RELMA® are registered United States trademarks of Regenstrief Institute, Inc.

Permission is hereby granted in perpetuity, without payment of license fees or royalties, to use, copy, or distribute the RELMA® program, RELMA® Users' Manual, RELMA® Release Notes, RELMA® database and associated search index files, LOINC® codes, LOINC® Users' Guide, LOINC® table (in all formats in which it is distributed by Regenstrief Institute, Inc. and the LOINC Committee), LOINC® Release Notes, LOINC® Changes File, LOINC® panels and forms file, and LOINC® hierarchies file (collectively, the "Licensed Materials") for any commercial or non-commercial purpose, subject to the following terms and conditions:

1) To prevent the dilution of the purpose of the LOINC codes and LOINC table of providing a definitive standard for identifying clinical information in electronic reports, users shall not use any of the Licensed Materials for the purpose of developing or promulgating a different standard for identifying patient observations, such as laboratory test results; other diagnostic service test results; clinical observations and measurements; reports produced by clinicians and diagnostic services about patients; panels, forms and collections that define aggregations of these observations; and orders for these entities in electronic reports and messages.

2) If the user elects to utilize the RELMA program, users receive the full RELMA database and associated search index files with the RELMA program, including the LOINC table and other database tables comprising the RELMA database. In addition to its use with the RELMA program, users may use the LOINC table by itself and may modify the LOINC table as permitted herein. Users may not use or modify the other database tables from the RELMA database or the associated search index files except in conjunction with their authorized use of the RELMA program, unless prior written permission is granted by the Regenstrief Institute, Inc. To request written permission, please contact [loinc@regenstrief.org](mailto:loinc@regenstrief.org).

3) Users shall not change the meaning of any of the LOINC codes. Users shall not change the name of, or any contents of, any fields in the LOINC table. Users may add new fields to the LOINC table to attach additional information to existing LOINC records. Users shall not change the content or structure of the LOINC panels and forms from the LOINC panels and forms file, but may notify the Regenstrief Institute of any potential inconsistencies or corrections needed by contacting [loinc@regenstrief.org](mailto:loinc@regenstrief.org).

4) A user may delete records from the LOINC table to deal with the user's local requirements. A user also may add new records to the LOINC table to deal with the users' local requirements, provided that if new records are added, any new entry in the LOINC\_NUM field of such new records must contain a leading alphabetic "X" so that the new codes and records cannot be confused with existing LOINC codes or new LOINC codes as they are defined in later releases of the LOINC table. Records deleted or added by users to deal with local requirements are not reflected in the official LOINC table maintained by the Regenstrief Institute and the LOINC Committee. Users must also make reasonable efforts to submit requests to LOINC for new records to cover observations that are not found in the LOINC table in order to minimize the need for X-codes.

5) LOINC codes and other information from the LOINC table may be used in electronic messages for laboratory test results and clinical observations such as HL7 ORU messages, without the need to include this Copyright Notice and License or a reference thereto in the message (and without the need to include all fields required by Section 7 hereof). When the LOINC code (from the LOINC\_NUM field) is included in the message, users are encouraged, but not required, to include the corresponding LOINC short name (from the SHORTNAME field) or the LOINC long common name (from the LONG\_COMMON\_NAME field) in the message if the message provides a place for a text name representation of the code.

6) Users may make and distribute an unlimited number of copies of the Licensed Materials. Each copy thereof must include this Copyright Notice and License, and must include the appropriate version number of the Licensed Materials if the Licensed Materials have a version number, or the release date if the Licensed Materials do not have a version number. This Copyright Notice and License must appear on every printed copy of the LOINC table. Where the Licensed Materials are distributed on a fixed storage medium (such as diskette or CD-ROM), a printed copy of this Copyright Notice and License must be included on or with the storage medium, and a text file containing this information also must be stored on the storage medium in a file called "license.txt". Where the Licensed Materials are distributed via the Internet, this Copyright Notice and License must be accessible on the same Internet page from which the Licensed Materials are available for download. This Copyright Notice and License must appear verbatim on every electronic or printed copy of the RELMA Users' Manual and the LOINC Users' Guide. The RELMA Users' Manual and the LOINC Users' Guide may not be modified, nor may derivative works of the RELMA Users' Manual or LOINC Users' Guide be created, without the prior written permission of the Regenstrief Institute, Inc. To request written permission, please contact [loinc@regenstrief.org](mailto:loinc@regenstrief.org). The Regenstrief Institute retains the right to approve any modification to, or derivative work of, the RELMA Users' Manual or the LOINC Users' Guide.

7) Subject to Section 1 and the other restrictions hereof, users may incorporate portions of the LOINC table, LOINC panels and forms file, and LOINC hierarchies file into another master term dictionary (e.g. laboratory test definition database), or software program for distribution outside of the user's corporation or organization, provided that any such master term dictionary or software program includes the following fields reproduced in their entirety from the LOINC table: LOINC\_NUM, COMPONENT, PROPERTY, TIME\_ASPCT, SYSTEM, SCALE\_TYP, METHOD\_TYP, STATUS, and SHORTNAME. Users are also required to either: (1) include the EXTERNAL\_COPYRIGHT\_NOTICE or (2) delete the rows that include

third party copyrighted content (e.g., third party survey instruments and answers). If third party content is included, users are required to comply with any such third party copyright license terms. Users are encouraged, but not required, to also include the RelatedNames2 and the LONG\_COMMON\_NAME in any such database. Further description of these fields is provided in Appendix A of the LOINC Users' Guide. Every copy of the LOINC table, LOINC panels and forms file, and/or LOINC hierarchies file incorporated into or distributed in conjunction with another database or software program must include the following notice:

"This product includes all or a portion of the LOINC® table, LOINC panels and forms file, and/or LOINC hierarchies file, or is derived from one or more of the foregoing, subject to a license from Regenstrief Institute, Inc. Your use of the LOINC table, LOINC codes, LOINC panels and forms file, and LOINC hierarchies file also is subject to this license, a copy of which is available at <http://loinc.org/terms-of-use>. The current complete LOINC table, LOINC Users' Guide, LOINC panels and forms file, and LOINC hierarchies file are available for download at <http://loinc.org>. The LOINC table and LOINC codes are copyright © 1995-2010, Regenstrief Institute, Inc. and the Logical Observation Identifiers Names and Codes (LOINC) Committee. The LOINC panels and forms file and LOINC hierarchies file are copyright © 1995-2010, Regenstrief Institute, Inc. All rights reserved. THE LOINC TABLE (IN ALL FORMATS), LOINC PANELS AND FORMS FILE, AND LOINC HIERARCHIES ARE PROVIDED "AS IS." ANY EXPRESS OR IMPLIED WARRANTIES ARE DISCLAIMED, INCLUDING, BUT NOT LIMITED TO, THE IMPLIED WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE. LOINC® is a registered United States trademark of Regenstrief Institute, Inc. A small portion of the LOINC table may include content (e.g., survey instruments) that is subject to copyrights owned by third parties. Such content has been mapped to LOINC terms under applicable copyright and terms of use. Notice of such third party copyright and license terms would need to be included if such content is included."

If the master term dictionary or software program containing the LOINC table, LOINC panels and forms file, and/or LOINC hierarchies file is distributed with a printed license, this statement must appear in the printed license. Where the master term dictionary or software program containing the LOINC table, LOINC panels and forms file, and/or LOINC hierarchies file is distributed on a fixed storage medium, a text file containing this information also must be stored on the storage medium in a file called "LOINC\_short\_license.txt". Where the master term dictionary or software program containing the LOINC table, LOINC panels and forms file, and/or LOINC hierarchies file is distributed via the Internet, this information must be accessible on the same Internet page from which the product is available for download.

8) Use and distribution of the Licensed Materials in ways that are not specifically discussed herein shall always be accompanied by the notice provided in Section 7 hereof. The guidelines for providing the notice that are contained in the last paragraph of Section 7 also shall apply. If a user has a question about whether a particular use of any of the Licensed Materials is permissible, the user is invited to contact the Regenstrief Institute by e-mail at [loinc@regenstrief.org](mailto:loinc@regenstrief.org).

9) If the user desires to translate any of the Licensed Materials into a language other than English, then user shall notify Regenstrief via email at [loinc@regenstrief.org](mailto:loinc@regenstrief.org). Any such

translation is a derivative work, and the user agrees and does hereby assign all right, title and interest in and to such derivative work: (1) to Regenstrief and the LOINC Committee if the translation is a derivative of the LOINC codes, LOINC Users' Guide, or LOINC table, and (2) to Regenstrief if the translation is a derivative work of the RELMA program, LOINC panels and forms file, LOINC hierarchies file, RELMA Users' Manual, RELMA database or associated search index files. Further, user shall fully cooperate with Regenstrief in the filing and reviewing of any copyright applications or other legal documents, and signing any documents (such as declarations, assignments, affidavits, and the like) that are reasonably necessary to the preparation of any such copyright application. The assignment granted by this paragraph extends to all proprietary rights both in the United States, and in all foreign countries. No other right to create a derivative work of any of the Licensed Materials is hereby granted (except the right to translate into a language other than English granted in this Section **Error! Reference source not found.**, and Regenstrief and the LOINC Committee respectively reserve all other rights not specifically granted herein. All such translations shall be electronically transmitted to Regenstrief, and such translations shall be made available and are subject to the same license rights and restrictions contained herein. Regenstrief will give credit on its website (and on screens in RELMA and in its users guides) to the user and/or entity that did the translation.

10) The Regenstrief Institute, Inc. and the LOINC Committee welcome requests for new LOINC content (terms, codes or associated material such as text descriptions and synonyms) and suggestions about revisions to existing content within the Licensed Materials. Any content submitted in conjunction with such a request is subject to the LOINC Submissions Policy, which is available at <http://loinc.org/submissions-policy>.

11) The names "Regenstrief," "Regenstrief Foundation," "Regenstrief Institute," and "LOINC Committee" may not be used in a way which could be interpreted as an endorsement or a promotion of any product or service without prior written permission of the Regenstrief Institute, Inc. Further, no right to use the trademarks of Regenstrief is licensed hereunder. To request written permission, please contact [loinc@regenstrief.org](mailto:loinc@regenstrief.org).

12) **DISCLAIMER: REGENSTRIEF INSTITUTE, INC. AND THE LOINC COMMITTEE, AS WELL AS ANY CONTRIBUTORS WHO HAVE PROVIDED TRANSLATIONS OF THE LICENSED MATERIALS, DO NOT ACCEPT LIABILITY FOR ANY OMISSIONS OR ERRORS IN THE LICENSED MATERIALS OR ANY OTHER MATERIALS OBTAINED FROM REGENSTRIEF INSTITUTE, INC. AND/OR THE LOINC COMMITTEE. THE LICENSED MATERIALS AND ALL OTHER MATERIALS OBTAINED FROM REGENSTRIEF INSTITUTE, INC. AND/OR THE LOINC COMMITTEE ARE PROVIDED "AS IS," WITHOUT WARRANTY OF ANY KIND. ANY EXPRESSED OR IMPLIED WARRANTIES ARE HEREBY DISCLAIMED, INCLUDING, BUT NOT LIMITED TO, THE IMPLIED WARRANTIES OF TITLE, NON-INFRINGEMENT, MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE AND WARRANTIES ARISING FROM A COURSE OF DEALING, TRADE USAGE, OR TRADE PRACTICE. FURTHER, NO WARRANTY OR REPRESENTATION IS MADE CONCERNING THE ACCURACY, COMPLETENESS, SEQUENCE, TIMELINESS OR AVAILABILITY OF THE LICENSED MATERIALS OR ANY OTHER MATERIALS OBTAINED FROM REGENSTRIEF INSTITUTE, INC. AND/OR THE LOINC COMMITTEE, OR ANY**

**TRANSLATIONS OR DERIVATIVE WORKS OF ANY OF THE FOREGOING. IN NO EVENT SHALL REGENSTRIEF INSTITUTE, INC. OR THE LOINC COMMITTEE OR ITS CONTRIBUTORS BE LIABLE FOR ANY DIRECT, INDIRECT, INCIDENTAL, SPECIAL, EXEMPLARY, RELIANCE, OR CONSEQUENTIAL DAMAGES OR ATTORNEYS' FEES (INCLUDING, BUT NOT LIMITED TO, PROCUREMENT OF SUBSTITUTE GOODS OR SERVICES; OPPORTUNITY COSTS; LOSS OF USE, DATA, SAVINGS OR PROFITS; OR BUSINESS INTERRUPTION) HOWEVER CAUSED AND ON ANY THEORY OF LIABILITY WHETHER IN CONTRACT, STRICT LIABILITY, OR TORT (INCLUDING NEGLIGENCE OR OTHERWISE) ARISING IN ANY WAY OUT OF THE USE OF THE LICENSED MATERIALS OR ANY OTHER MATERIALS OBTAINED FROM REGENSTRIEF INSTITUTE, INC. AND/OR THE LOINC COMMITTEE, EVEN IF ADVISED OF THE POSSIBILITY OF SUCH DAMAGE OR IF SUCH DAMAGES WERE FORESEEABLE. SOME JURISDICTIONS DO NOT ALLOW THE LIMITATION OR EXCLUSION OF CERTAIN WARRANTIES OR CONDITIONS, SO SOME OF THE FOREGOING MAY NOT APPLY TO YOU.**

13) This license shall be construed and interpreted in accordance with the laws of the State of Indiana, United States of America, excluding its conflicts of law rules.

#### Notice of Third Party Content and Copyright Terms

A small portion of the content of the LOINC table, LOINC panels and forms, LOINC hierarchies, RELMA database and associated search index files consists of content subject to copyright from third parties. This third party content is either used with permission or under the applicable terms of use. In all such cases, we have included the copyright notice. This third party content is highlighted in the program as follows: When such copyright content appears in the RELMA look-up grid, RELMA will highlight the row containing that content by printing in a different background color and using italics. It will also include a link in the (EXT (C)) column. By clicking on that link, users will get to the copyright notice and to the terms of use for the content of those LOINC-mapped terms. In the case of a LOINC table (e.g. the tab delimited file or the LOINC Access database) we include the copyright notice (up to 250 characters).

We have included third party content that allows use and distribution at least for clinical, administrative and research purposes. The third party copyright owners generally ask for attribution of the source, allow the free use of the content for treatment, health care management, and research purposes. They generally forbid alteration of their content (e.g., survey questions and/or answers) and use for commercial purpose, which usually means the direct sale of the survey instruments, but they often do allow use of their content in commercial software, medical record and other clinical database systems, and the messaging of patient information collected through the use of these instruments. The details of the notice of copyright for any third party content can be found in association with the terms when using the RELMA look-up tool or the LOINC table. The copyright of the LOINC codes per se remain owned by Regenstrief Institute, Inc. and the LOINC Committee and subject to the LOINC Copyright Notice and License.

In the future, we expect to include many more survey instruments and questionnaires from third parties with permission (especially those required by the U.S. federal government for payment and reimbursement) and believe that cataloguing all of these data collection forms in one comprehensive system (the LOINC table) along with laboratory and other clinical variables will facilitate the use of this

data in direct clinical care, research and practice management.

## Preface

The LOINC database provides a set of universal names and ID codes for identifying laboratory and clinical test results.<sup>1,2</sup> LOINC facilitates the exchange and pooling of results, such as blood hemoglobin, serum potassium, or vital signs, for clinical care, outcomes management, and research. Currently, many laboratories use ASTM 1238<sup>3</sup> or its sister standard, HL7<sup>4</sup>, to send laboratory results electronically from production laboratories to clinical care systems in hospitals. Most laboratories identify tests in HL7 messages by means of their internal (and idiosyncratic) code values. Receiving medical informatics systems cannot fully “understand” the results they receive unless they either adopt the producer's laboratory codes (which is impossible if informatics system receives results from multiple source laboratories, e.g., the hospital lab, the local commercial lab, and a nursing home lab), or invest in the work to map each laboratory's coding system to their internal code system.<sup>5</sup>

If medical information producers who wish to communicate with each other adopt LOINC codes to identify their results in data transmissions, this problem would disappear. The receiving system with LOINC codes in its master vocabulary file would be able to understand and properly file HL7 results messages that identified clinical observations via LOINC codes. Similarly, if test and observation codes were reported test with the LOINC codes, government agencies would be able to pool results for tests from many sites for research management and public health purpose. The LOINC codes (and names) for test observations should be of interest to hospitals, clinical laboratories, doctors' offices, state health departments, governmental health care providers, third-party payers, and organizations responsible for quality assurance and utilization review.

The LOINC codes are not intended to transmit all possible information about a test or observation. They are only intended to identify the test result or clinical observation. Other fields in the message can transmit the identity of the source laboratory and special details about the sample. (For instance, the result code may identify a blood culture, but the message source code can be more specific and identify the sample as pump blood.) The level of detail in the LOINC definitions was intended to distinguish tests that are usually distinguished as separate test results within the master file of existing laboratory systems. Indeed, at the outset, we used the master files from seven U.S. laboratories to shape this effort, and requests from commercial labs and hospitals continue to shape the content of the LOINC effort.

Each LOINC record corresponds to a single test result or panel. The record includes fields for specifying:

1. Component (analyte) - e.g., potassium, hemoglobin, hepatitis C antigen.
2. Property measured - e.g., a mass concentration, enzyme activity (catalytic rate).
3. Timing - i.e., whether the measurement is an observation at a moment of time, or an observation integrated over an extended duration of time - e.g., 24-hour urine.
4. The type of sample - e.g., urine, blood.
5. The type of scale - e.g., whether the measurement is quantitative (a true measurement) ordinal (a ranked set of options), nominal (e.g., *E. coli*; *Staphylococcus aureus*), or narrative (e.g., dictation results from x-rays).
6. Where relevant, the method used to produce the result or other observation.

It also contains information about the amount, route, and timing of physiologic or pharmacologic challenges (e.g., oral glucose tolerance test, which would be expressed in LOINC as GLUCOSE^1H POST 100 G GLUCOSE PO1). The LOINC identifiers do not usually include the method in the name for chemistry tests, where tests are more often standardized to normalized methods; they do include methods for most serological tests and coagulation studies. This same principle is usually reflected in the master

files of existing laboratories. Of course, the method can always be reported as a separate item of information in a result message regardless of whether it is part of the test name.

We used many sources for constructing the database, including the Silver Book from the International Union of Pure and Applied Chemistry (IUPAC) and the International Federation of Clinical Chemistry (IFCC),<sup>6</sup> textbooks of clinical pathology (e.g., Henry<sup>7</sup> and Tietz<sup>8</sup>), the expertise and work of the LOINC members, and EUCLIDES. We have also reviewed the master test files of seven sources (Indiana University/Regenstrief, University of Utah, Association of Regional and University Pathologists (ARUP), Mayo Medical Laboratories, LDS Hospital in Salt Lake City, the Department of Veterans Affairs, Quest Diagnostics, and University of Washington). This has been an empirical effort. Our goal is to provide codes that correspond to the concepts in real world laboratories' and clinical departments' master files.

The database includes fields for each of the six parts of the name. In addition, it also contains short names (as of the August 2002 version for laboratory tests), related words, synonyms, and comments for all observations. Related words (synonyms) are included to facilitate searches for individual laboratory test and clinical observation results.

We have defined fields in the database for a number of data elements, e.g., typical units, sample normal ranges, but most of those fields are only partially populated. In a few cases, we have suggested standard answer lists for tests whose results are usually reported as codes. The database is an ongoing project. We have established guidelines for users who wish to request additions and changes to LOINC, which are detailed in Appendix D.

For some kind of tests and observations, the database provides several ways to report values. For example, blood cell antigens might be presented as a "panel" with separate "tests" which report each possible antigen as present or absent if the test is to establish paternity; for cross matching, the result would be reported as a list of antigens found. We try to provide for both methods of reporting in the LOINC databases by including codes for both types of test identifiers.

Laboratories and managers of medical records systems should record the LOINC codes as attributes of their existing test/observation master files and use the LOINC codes and names in the OBSERVATION ID field (OBX-3) of the ASTM and HL7 OBX segment and the corresponding CEN TC251 and DICOM messages to identify laboratory results.

The overall organization of the database is divided first into four categories, "lab", "clinical", "attachments" and "surveys". (This split is recorded in CLASSTYPE.) The laboratory portion is further divided into the usual categories of chemistry, hematology, serology, microbiology (which includes parasitology and virology), and toxicology. We have separated antibiotic susceptibilities into their own category. The clinical portion of the LOINC database contains entries for vital signs, hemodynamics, intake/output, EKG, obstetric ultrasound, cardiac echo, urologic imaging, gastroendoscopic procedures, pulmonary ventilator management, and other clinical observations. Appendix B lists these classes in more detail. There is nothing sacred about these categories, and you are free to sort the database by whatever class is convenient for your application.

The Regenstrief Institute maintains the LOINC database and makes it available in a number of file formats. In each of them, the first part of the file contains the copyright notice with permission to use the database for any purpose without charge or written permission. We have copyrighted the databases and this document to assure that multiple variants of the codes do not emerge. Having many variants would defeat the purpose of a universal identifier for test results. The LOINC database (which identifies over 34,000 different lab tests and clinical observations), supporting documentation and the RELMA<sup>®</sup> mapping program are all available through the Regenstrief Institute web site. (<http://loinc.org>)

**LOINC ACCESS database:**

The official LOINC database is available as an ACCESS file called LOINC.MDB. It was created using Microsoft Access<sup>™</sup> 2007.

**LOINC Tab Delimited ASCII:**

Each record of the database is on a separate line. Each record is terminated by CR/LF, and each field is delimited with a tab character. Non-null text fields are enclosed in double quotes (“). This is the format you will use if you want to import into your own database. This file contains all of the content of the database and is formatted to be easily imported into a wide variety of database and spreadsheet applications.

**The LOINC Users' Guide** (this document) is available as a PDF file. It explains the structure of the database, its rationale, and the rules we used for naming test results.

**RELMA**

In addition to the basic LOINC files, we produce a Windows-based mapping utility called the Regenstrief LOINC Mapping Assistant (RELMA<sup>®</sup>). This program is also available for free use

The RELMA package includes the LOINC table in the database plus several large index tables.

**RELMA Users' Manual**

There is a separate Users' Manual documenting the RELMA program.

All of the above files are available from the LOINC website <http://loinc.org> . They are also distributed on CD.

We welcome corrections or extensions to the database. We are not interested in adding terms that might be needed in some future situation but we are interested in adding test observations that are actively being reported today. Appendix D provides instructions for submitting new terms.

Clem McDonald  
Chairman, LOINC Committee  
Chairman, Laboratory LOINC Committee

Stan Huff  
Co-Chairman, LOINC Committee  
Chairman, Clinical LOINC Committee

**Acknowledgments**

We wish to thank Henrik Olesen, Chairman of IUPAC, Commission on Quantities & Units in Clinical Chemistry, for his very helpful comments and insights about laboratory test coding.

This endeavor was supported in part by Grant Numbers R13/CCR517099 and H75/CCH522778 from the Centers for Disease Control and Prevention. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the Centers for Disease Control and Prevention. This work also was supported in part by grants and contracts from the John A. Hartford Foundation of New York, the National Library of Medicine (Contracts N01-LM-4-3510, N01-LM-6-3546, N01-LM-9-3517 and N01-LM-3-3501), and the Agency for Health Care Policy and Research (AHCPR) (Grants HS 08750 and HS 07719 ). This work was initiated by and performed under the auspices of the Regenstrief Institute.

# 1 Introduction

The goal of the LOINC project is to create universal identifiers (names and codes) used in the context of existing ASTM E1238, HL7, CEN TC251, and DICOM observation report messages employed in the various sub-domains of healthcare informatics such as Clinical Laboratory Information Management Systems and Computer-Based Patient Record Systems.<sup>9, 10</sup> Specifically, the identifiers can be used as the coded value of the "Observation Identifier" field (# 3) of the OBX segment of an ORU HL7 (HL7 vs. 2.x and vs. 3.9 or ASTM 1238-9410) messages, or in a corresponding field in future versions of these HL7 and DICOM standards. LOINC codes identified in HL7 as code system "LN" provide "universal" identifiers. When used in the context of the messaging standards, LOINC codes allow the exchange of clinical laboratory data between heterogeneous computing environments.

To facilitate this process, each identifier requires a fully specified name created in a standard way so that users can create long names for their tests that can be linked to the universal test identifier using semi-automated methods.

We focused our initial effort on creating names for results of reportable tests or clinical measurements rather than request-able batteries, because the issues involved in naming results of tests are less complex than those involved in naming the batteries. However, we have also defined codes for some order panels. It is important to note that LOINC codes for single tests, reports, and observations are equally suitable for the ordered item in an order record or message, or as the result identifier in a result message.

The LOINC database is a "universal" master file of standard "test" names and codes that will cover most of the entries in these files of operational laboratory systems, so that the terms in these operational master files could be mapped directly to universal codes and names. The names we create correspond most closely to the "long test descriptions" seen in test master files. The LOINC names are "fully specified" names. That is, if a person wanted to map her local test dictionary to the LOINC codes, all the information needed to map a local test name to one of the fully specified names should be present in the LOINC name.

We aim to achieve a level of detail in the definition of a test that will map one-to-one to the separately reported observations on a clinical laboratory report. If a test has its own column on a clinical report, or has a reference range that is significantly different from other tests, or has a different clinical meaning than other related tests, it will usually be assigned a separate LOINC code and name. We deliver these fully specified names, their codes, and their related names as a database in which each line corresponds to a unique test measurement.

## *1.1 Successes*

The LOINC codes have been greeted enthusiastically since they were released to the Internet in April of 1996. Since then we have released thirteen revisions of the LOINC database and it now includes over 30,000 observation concepts. The informatics committee of the College of American Pathologists (CAP) has endorsed the LOINC codes. The American Clinical Laboratory Association (ACLA), an association of large referral laboratories whose members are responsible for more than 60% of US outpatient laboratory test volume, has recommended LOINC for adoption by its members. Quest Diagnostics<sup>®</sup> (formerly Corning MetPath), LabCorp<sup>®</sup>, and SmithKline Beecham (now part of Quest Diagnostics), three of the largest commercial laboratories in the US, have adopted LOINC as their code system for reportable test results, as has ARUP (Associated Regional and University Pathologists). Mayo Medical Laboratories

is currently mapping their tests to LOINC. In addition, the University of Colorado, Intermountain Health Care, Kaiser Permanente<sup>®</sup>, Clarian Health (Indiana University, Methodist Hospital, and Riley Hospital), Partners Healthcare System of Boston<sup>™</sup> (Brigham and Women's and Mass General Hospital), Care Group of Boston<sup>®</sup>, Mayo Clinic, and the Department of Defense are adopting the LOINC codes for laboratory reporting. All US veterinary medicine laboratories have committed to the use of LOINC.

HMOs such as Empire Blue Cross<sup>®</sup> and Aetna<sup>®</sup> Health Care are also adopting LOINC for internal purposes. Internationally, LOINC has also met success. Geneva, Switzerland, is adopting LOINC for quality assurance mandates. The provinces of Ontario and British Columbia, Canada, are adopting LOINC codes province wide, and Newfoundland is considering following in their footsteps. Most recently, Germany has adopted LOINC for national use.

The LOINC codes have been incorporated into the National Library of Medicine's Unified Medical Language System<sup>®</sup> (UMLS<sup>®</sup>). They have been incorporated in HCFA's quality assurance testing pilot programs, and part of the draft Health Insurance Portability and Accountability Act (HIPPA) electronic attachments specification. They have been adopted by the Centers for Disease Control and Prevention/Council of State and Territorial Epidemiologists' project for electronically reporting/transmitting communicable disease information<sup>11,12</sup> and by the North American Association of Central Cancer Registries (NAACCR) for their tumor registry variables.

On March 21, 2003, the United States Departments of Health and Human Services (HHS), Defense (DoD) and Veterans Affairs (VA) announced the first set of uniform standards for the electronic exchange of clinical health information to be adopted across the federal government. As part of this, all federal agencies that deal with health care data will adopt laboratory Logical Observation Identifiers Name Codes (LOINC) to standardize the electronic exchange of clinical laboratory orders and results.

## ***1.2 What is not part of the name***

Certain parameters and descriptions pertaining to test performance are specifically excluded from the fully specified test name. These parameters will typically be reported in separate fields (attributes) of a test/observation report message, not as part of the observation name. Attributes that we explicitly exclude from the fully specified name are:

- the instrument used in testing
- fine details about the sample or the site of collection such as “right antecubital fossa”
- the priority of the testing, e.g., whether stat or routine
- who verified the result
- the size of the sample collected
- the place of testing (e.g., home, bedside, clinical lab)

In the case of laboratory tests, the name does include information that identifies the type of sample (or specimen). However, the “sample” part of the name is not meant to carry all possible information about the sample, but only enough to indicate significant differences in the result and to reflect current usage in test names. For example, laboratories usually define urine sodium, sweat sodium, and serum sodium as different tests because each of these has a different normal range. But laboratories do not define different tests to distinguish the concentration of arterial serum sodium from venous serum sodium, though the lab may report that the sample was venous or arterial in another part of the report. We are guided by the pragmatics of conventional usage. If laboratories define separate tests for the same measurements done on different specimens (this usually implies a well-defined normal range difference), we will define

different “result-able” tests in our dictionary. If they do not, we will not.

The extent to which we include methods as part of the name is also guided by pragmatics. We distinguish tests/observations by the type of method used to produce the results only if a given type of method has an important effect on the interpretation of the result. This is a complex subject and it is difficult to fully describe our rationale in this report. Where laboratories do not tend to include the method in the name (e.g., most of chemistry) we do not include the method in the name. Where they tend to (e.g., in immunochemistry) we do. For some tests, this can be justified by the standardization of methods to produce “equivalent” results, and sometimes by the many variables (method, reagent) that one could never hope to represent fully in a single name. However, even when we do distinguish these cases, we distinguish by type of method, not the most detailed possible method distinction. (See section 2.7, Type of Method, for more details.)

The College of American Pathologists produces statistical summaries of the results for measurements of standard samples broken down by laboratory and by instrument or procedure. (These are called CAP surveys.) We considered using this CAP survey data to decide empirically when test names should be distinguished by method, but decided this was not feasible because many of the apparent differences in method obtained with the standard samples were artifacts of the sample matrix and did not apply to serum specimens. In addition, the variation among laboratories was often of the same magnitude as the variation among methods within laboratories for the same method.

We do not mean to underrate the importance of method differences. The result message will still include information about the normal range for that particular test, the source laboratory and, if the laboratory wishes, specific information about the method (e.g., OBX 17 can carry very specific method information). However, such information is reported in separate fields in the HL7 message. It is not embedded in the names of the test.

### *1.3 Scope of LOINC*

The current scope of the existing laboratory portion of the LOINC database includes all observations reported by clinical laboratories, including the specialty areas: chemistry, including therapeutic drug monitoring and toxicology; hematology; serology; blood bank; microbiology; cytology; surgical pathology; and fertility. A large number of terms used in veterinary medicine have also been included. In addition, the scope includes those non-test measurements that are commonly required to interpret test results and are usually included as part of the report with the laboratory observations. Examples include:

- for cervical pap smears, the phase of menstrual cycle or use of estrogens
- for arterial blood gases, inspired oxygen
- for drug concentrations used in pharmacokinetics, the dose
- for a blood bank, the number of units dispensed

The June 2000 release contained our first foray into order sets/batteries. Existing LOINC codes could always be used to order the specific tests observation, but prior to 2000 there was no mechanism to use LOINC codes to order a set of observations. We have currently only addressed a group of observations that are either naturally produced as a panel (e.g., urinalysis) or are defined by some national body (e.g., Basic metabolic HCFA 2000 panel).

The clinical portion of the LOINC database covers the areas of blood pressure, heart and respiratory rates, critical care measures, cardiac output, body dimensions, body temperature, intake and output, electrocardiography, cardiac echo, obstetric ultrasound, urologic ultrasound, gastrointestinal endoscopy,

ventilator management, dental, Data Elements for Emergency Department Systems (DEEDS) reporting, radiology study reporting, claims attachment and the major headings of history and physical, discharge summary, and operative note reports and tumor registry variables. Further work on clinical obstetrics and nursing observations is ongoing. There are separate sections for Claims Attachments and Survey Instruments.

### 1.4 The LOINC Code Identifier

To each name, we have assigned a unique permanent code that we call the LOINC code. This is the code that systems should use to identify test results in electronic reports. The LOINC code has no intrinsic structure except that the last character in the code is a mod 10-check digit. The algorithm to calculate this check digit is given in Appendix C. All of the structure associated with a single LOINC entity is stored in other fields in the LOINC database.

## 2 Major “Parts” of a Test/Observation Name

The fully specified name of a test result or clinical observation has five or six main parts including: the name of the component or analyte measured (e.g., glucose, propranolol), the property observed (e.g., substance concentration, mass, volume), the timing of the measurement (e.g., is it over time or momentary), the type of sample (e.g., urine, serum), the scale of measurement (e.g., qualitative vs. quantitative), and where relevant, the method of the measurement (e.g., radioimmunoassay, immune blot). These can be described formally with the following syntax.

**<Analyte/component>:<kind of property of observation or measurement>:<time aspect>:<system (sample)>:<scale>:<method>**

The colon character, “:”, is part of the name and is used to separate the main parts of the name.

The first part of the name can be further divided up into three subparts, separated by carats (^). The first subpart can contain multiple levels of increasing taxonomic specification, separated by dots (.). The third and fourth parts of the name (time aspect and system/sample) can also be modified by a second subpart, separated from the first by a carat. In the case of time aspect, the modifier can indicate that the observation is one selected on the basis of the named criterion (maximum, minimum, mean, etc.); in the case of system, the modifier identifies the origin of the specimen if not the patient (e.g., blood donor, fetus, and blood product unit). The hierarchical structure is outlined in Table 1, with references to the section numbers where each item is explained in detail.

<b>Subpart Name</b>	<b>Section</b>
Component/analyte	2.2
Name and modifier	2.2.1
Component/analyte name	2.2.1.1
Component/analyte subname	2.2.1.2
Component/analyte sub-sub-name	2.2.1.3
Information about the challenge (e.g., 1H post 100 gm PO challenge)	2.2.2
Adjustments/corrections	2.2.3
Kind of Property (mass concentration, mass)	2.3

Time Aspect (point or moment in time vs. time interval)	2.4
System/Sample type (urine, serum)	2.5.1
“Super System” (patient, donor, blood product unit)	2.5
Type of Scale (nominal, ordinal, quantitative)	2.6
Method Type	2.7

We used Tietz<sup>13</sup>, Henry<sup>14</sup>, IUPAC<sup>15</sup>, EUCLIDES<sup>16</sup>, diagnostic microbiology textbooks, such as Mahon and Manuselis<sup>17</sup>, the American Association of Blood Banking<sup>18</sup>, and other sources as well as the expertise of the individuals or the committee to choose preferred names.

Examples of fully specified LOINC names:

Sodium:SCnc:Pt:Ser/Plas:Qn  
Sodium:SCnc:Pt:Urine:Qn  
Sodium:SRat:24H:Urine:Qn  
Creatinine renal clearance:VRat:24H:Ur+Ser/Plas:Qn  
Glucose^2H post 100 g glucose PO:MCnc:Pt:Ser/Plas:Qn  
Gentamicin^trough:MCnc:Pt:Ser/Plas:Qn  
ABO group:Type:Pt:Bld^donor:Nom  
Body temperature:Temp:8H^max:XXX:Qn  
Chief complaint:Find:Pt:^Patient:Nar:Reported  
Physical findings:Find:Pt:Abdomen:Nar:Observed  
Binocular distance:Len:Pt:Head^fetus:Qn:US.measured

## 2.1 General naming conventions

### 2.1.1 Abbreviations in names of component/analyte

Except for enumerated exceptions (Table 2), abbreviations should not be used in the component (analyte) of the name. We require the use of “total”, not “tot”, “fraction”, not “frac”, “Alpha”, not “A-,” “Beta” not “B-” (and so on for any Greek letter), “oxygen”, “not O<sub>2</sub>”, and so on.

<b>Abbreviation</b>	<b>Full Name</b>
Ab	Antibody
Ag	Antigen
DNA	deoxyribonucleic acid
HIV	human immunodeficiency virus
HLA	human histocompatibility complex derived antigens

HTLV 1	human t-cell lymphotropic virus-1
Ig "X"	immunoglobulins (e.g., IgG for immunoglobulin G, IgM for immunoglobulin M)
NOS	not otherwise specified
RNA	ribonucleic acid
rRNA	ribosomal ribonucleic acid

- 2.1.2 General naming rules for the component (analyte) part of the fully specified name.
- 2.1.2.1 Place the identifier of the substance being measured first. This means "Hepatitis A antibodies (Ab)" not "Antibodies, Hepatitis A."
- 2.1.2.2 **Use the generic name of a drug**, not the brand name, when referring to drug concentrations and antimicrobial susceptibilities, e.g., Propranolol, not Inderal. We will usually include the brand or trade names in the related names (synonyms) field.
- 2.1.2.3 Use full taxonomic name of an organism or virus name (not the disease) when describing a test that diagnoses that disease. Say "Rickettsia rickettsii Ab" not "Rocky Mountain spotted fever Ab". Say "herpes simplex virus Ab" not "HSV Ab." The disease name should be included as a synonym in the related name field.
- 2.1.2.4 **Species and groups of species:** SP identifies a single species whose identity is not known. SPP identifies the set of species beneath a genus. We have a third case, however. In some tests, antibodies apply to different strains of species. In rickettsial diseases, the antibodies are then against groups of species, e.g., the spotted fever group or the typhus group. In this case we use Rickettsia spotted fever group and Rickettsia typhus group.
- 2.1.2.5 When tests include the name of a bacterium (e.g., Neisseria gonorrhoeae DNA probe) for the formal LOINC name we use the full bacterial name from the International Journal of Systematic and Evolutionary Microbiology<sup>19</sup>. When it includes the name of a virus (e.g., West Nile Virus IgM antibodies), we use the viral name as given by Index Virum<sup>20</sup>.
- 2.1.2.6 When the test measures an antigen to a specific species of organism but cross-reactivity is such that other organisms are identified, the name should be the principal organism that is targeted by the test.
- 2.1.2.7 Avoid "direct" and "indirect" except as parts of synonym names. Avoid "conjugated" and "unconjugated" when a more precise term, such as "glucuronidated" or "albumin-bound" is available.
- 2.1.2.8 Use "platelets", not "thrombocytes."
- 2.1.2.9 **Name vitamins by the chemical name.** For example, use thiamine not Vitamin B1, The name containing "Vitamin" will be included as a synonym. This is the only reasonable approach because all vitamins have a chemical name but not all vitamins have a "numbered" vitamin name.
- 2.1.2.10 **Always specify whether serology tests measure the antigen or antibody**, using the abbreviation "Ab" for antibody and "Ag" for antigen. Remove the "anti" from "ANTI X Ab." It is redundant and obscures the most significant word in the name. Thus, "anti-smooth muscle Ab" becomes "Smooth muscle Ab." Common abbreviations or shortened names, e.g., ANA for

anti-nuclear antibody, will be found in the related names field.

- 2.1.2.11 VDRL will be named Reagin Ab because that is what it is. We will have to depend upon synonyms and aliases to equate our “standardized” names with the old names.
- 2.1.2.12 Use the noun form of the target of the antibody, e.g., Myocardium Ab, not Myocardial Ab.
- 2.1.2.13 **Anion vs. acid:** Always use the anionic name for chemicals, not the acid name, e.g., lactate, citrate, and urate, not lactic acid, citric acid, and uric acid. The acid form of the name will be included in the related names field of the database.
- 2.1.2.14 **Alcohols:** Always use the single-word names for alcohols: methanol, not methyl alcohol; ethanol, not ethyl alcohol, and so on.
- 2.1.2.15 Always spell out OH as Hydroxy, or as - ol, with no space or hyphen between Hydroxy and the next word.
- 2.1.2.16 Greek letters, alpha, beta, gamma, etc., are always spelled out (e.g., alpha tocopherol, not A-tocopherol), with a space between the spelled out Greek letter and the rest of the chemical name
- 2.1.2.17 Use pH, not log (H<sup>+</sup>).
- 2.1.2.18 Whenever possible, the component will contain the scientific names of allergens. NOTE: This is a new convention implemented in January 2002.
- 2.1.2.19 Avoid use of the word “total” in laboratory test names, except when denoting the denominator of a fraction. Thus it is Alkaline phosphatase, not Alkaline phosphatase.total, but Alkaline phosphatase.bone/Alkaline phosphatase.total.
- 2.1.2.20 For drug metabolites, we will use the “nor” form rather than “desmethyl”, e.g., for instance nordoxepin not desmethyldoxepin.

### 2.1.3 Punctuation in analyte names

A number of analyte names include punctuation characters such as commas, for example, to identify the position of multiple alkyl groups in a carbon chain. We will avoid special characters, e.g., commas, dashes, and parentheses, except where they are included in the name specified by IUPAC, the Chemical Abstract Service (CAS) convention, or another international convention. So commas will appear in multiple substitutions of alkyl chains per the CAS standard, dashes will appear in HLA antigen names, and parentheses (i.e., round brackets) will appear in the names of red blood cell antigens.

### 2.1.4 Case insensitivity

All names are case insensitive. Prior to December 2006, we used upper case in the database and our examples, but change to mixed case for easier readability. In electronic messages senders and receivers can use upper, lower or mixed case. However, the meanings should not be sensitive to case conversions to avoid any possibility of confusion when the information is sent over networks that may apply case conversion. To identify parts of the few names that by international convention are case sensitive, such as red blood cell antigens, we use the word “little” in front of the letter that is lower case. We use a similar

convention to indicate superscripts with the word SUPER. See examples in Table 3.

<b>Table 3: Example Case Specifying Conventions</b>	
<b>Our conventions</b>	<b>Standard mixed case</b>
L little u super little a	Lu <sup>a</sup>
little i-1 subtype	i-1 Subtype

### 2.1.5 Roman numerals vs. Arabic numerals

Whenever possible, numerals shall be represented in their Arabic form. However, when the conventional name uses Roman numerals as is the case for clotting factors such as factor VIII, the LOINC primary name will use Roman numerals and we define a synonym containing Arabic numerals.

## 2.2 Component/analyte (1st part)

The first main part consists of three subparts: (1) the principal name (e.g., the name of the analyte or the measurement); (2) the challenge or provocation, if relevant, including the time delay, substance of challenge, amount administered, and route of administration; and (3) any standardization or adjustment.

The three subparts of the first part follow this syntax:

```
<[analyte].[subclass].[sub-subclass]> ^
<[time delay] post [amount] [substance] [route]> ^
<adjustment>
```

In the above syntax, the carat (^) is a required delimiter and the "dot" (.) separates the analyte name from its subspecies.

This convention also implies that dots (.) and carats (^) cannot be a formal part of any of the words that are connected by these delimiters.

These subparts are described in greater detail below, Sections 2.2.1 through 2.2.3.

### 2.2.1 Analyte Name (1st subpart)

The first subpart names the analyte, including any relevant sub-classifications, separated from the main analyte name by dots.

#### 2.2.1.1 Analyte/Subclass

The principal name (the first subpart) can be divided further by subclass (e.g., Calcium by itself is one component, Calcium.ionized names another test that measures a subclass of calcium.) Subclasses are separated by dots. Examples of common subclasses include: bound, free, and bioavailable; ionized and non-ionized; glycosylated; glucuronidated and non-glucuronidated; IgA, IgD, IgE, IgG, and IgM as modifiers indicating the subspecies of antibodies. Note that bio-available is distinguished from free by including both free and partially bound moieties.

If the antibody is from a particular subclass of antibodies specify the subclass (IgM, IgG, IgA, or IgD) e.g., Hepatitis A virus Ab.IgG, Hepatitis A virus Ab.IgM

If more than one species is included in the measurement, all are listed in the subclass, e.g., “Mumps virus Ab.IgG+IgM” with a plus sign (+) to separate the subspecies. There should be no spaces between the plus sign and the words it connects. If two constituents are measured as one quantity, both should be named and the component separated by a plus sign (+), e.g., Cyclosporine+Metabolites.

If analytes are measured separately, such as in a panel, the analytes are separated by an ampersand (&) surrounded by spaces, e.g., ABO & Rh panels. In panels, each analyte is measured individually. Impressions provide another use of the ampersand (&), for example, Hepatitis A virus Ab.IgM & total impression. In the case of the Hepatitis antibody impression, both the IgM antibody and the total impression are described separately.

### 2.2.2 Challenge test (2nd subpart)

The second subpart contains information necessary to interpret “challenge” (or loading or tolerance) tests. Variables that report the result of a measurement taken a certain amount of time post challenge (e.g., glucose after an oral glucose tolerance test) must be distinguished according to the challenge and the time post challenge. Thus, the second subpart has a substructure that identifies the time interval or time difference and the challenge, using the following syntax, where the word “post” (or base line) is required.

<time delay> “post” <challenge>

where the challenge can be further characterized as

<amount given> <substance/treatment given> <route given>

An example of a challenge that used all parts would be: Aldosterone^1H post 25 mg captopril PO  
The time difference follows the syntax: n<S|M|H|D|W> where n is a number (possibly a decimal); S denotes seconds; M denotes minutes; H denotes hours; D denotes days; and W denotes weeks. The time delay can be preceded by a 'greater than' (>) sign, e.g., >4H. Table 4 lists some possible values for time difference, but any time specification that follows the above syntax would be legal.

In addition to specifying a time elapsed since challenge, the time delay slot can be used to name a clock time when the measurement was taken, e.g., Glucose^10 AM specimen, or to specify the ordering of specimens, e.g., ^1st specimen, ^2nd specimen. Use this syntax to indicate pre- and post-immunization specimens, acute and convalescent specimens, or a series of specimens for which no more detailed information is available.

Table 4: Example Time Delay Post Challenge			
BS	Baseline (time just before the challenge)		
PEAK	The time post drug dose at which the highest drug level is reached (differs by drug)		
TROUGH	The time post drug dose at which the lowest drug level is reached (varies with drug)		
RANDOM	Time from the challenge, or dose not specified (random)		
n minutes/hours/days/weeks/months/etc. after challenge begun: □			
1M	1 minute post challenge	6H	6 hours post challenge
2M	2 minutes post challenge	7H	7 hours post challenge
3M	3 minutes post challenge	8H	8 hours post challenge

4M	4 minutes post challenge	8H SHIFT	8 hours aligned on nursing shifts
5M	5 minutes post challenge	12H	12 hours post challenge
6M	6 minutes post challenge	24H	24 hours post challenge
7M	7 minutes post challenge	2D	2 days
8M	8 minutes post challenge	3D	3 days
9M	9 minutes post challenge	4D	4 days
10M	10 minutes post challenge	5D	5 days
15M	15 minutes post challenge	6D	6 days
20M	20 minutes post challenge	7D	7 days
25M	25 minutes post challenge	1W	1 week
30M	30 minutes post challenge	10D	10 days
1H	1 hour post challenge	2W	2 weeks
1.5H	1½ hour (90 min) post challenge	3W	3 weeks
2H	2 hours post challenge	4W	4 weeks
2.5H	2½hours post challenge	1MO	1 month (30 days) post challenge
3H	3 hours post challenge	2MO	2 months (60 days) post challenge
4H	4 hours post challenge	3MO	3 months (90 days) post challenge
5H	5 hours post challenge		

The second subpart is also used to describe measurements taken at a specified point after the beginning of an ongoing treatment, such as peritoneal dialysis, e.g., Creatinine^12H post peritoneal dialysis. More generally, this syntax can be used to indicate that observations were recorded, e.g., ^post partum, ^post surgery, or ^post EDTA therapy.

The syntax of the second subpart can be specified in various ways to indicate challenges of greater or lesser specificity, corresponding to the amount of detail the lab knows about the challenge specimen. Examples of the range of possibilities include:

Analyte	“^”	Time	“Post”	Amount	Sub/Treat	Route
11-Deoxycortisol	^	8H	post	30 mg/kg	Metyrapone	PO
Corticotropin	^	45M	post	dose u/kg	Insulin	IV
Ascorbate	^		post	dose		PO
11-Deoxycortisol	^	2 <sup>ND</sup> specimen	post		XXX challenge	
17-Hydroxyprogesterone	^	6H	post		XXX challenge	
11-Deoxycortisol	^		post		XXX challenge	
Calcium	^	12H	post		CFst	
C peptide	^		post		CFst	

### 2.2.2.1 Reporting the baseline measure as part of a challenge test

We define one baseline term for different challenge batteries when the challenge is given by the same

dose and route. So we define one baseline test for the 100 gm oral glucose tolerance test regardless of the number of separate measurements defined in the battery. For example, the baseline serum glucose for 100 gm oral glucose by mouth would be:

Glucose<sup>^</sup>pre 100 g glucose PO

A laboratory could use this same test identifier to identify the baseline result of a two hour glucose tolerance and a three hour glucose tolerance, for example.

We would define different baseline measurements for challenges with different substances. The baseline serum glucose before a challenge with 50 U insulin challenges would be defined as a different test from the baseline glucose for an oral glucose tolerance test. These different baseline tests are defined to accommodate laboratories that conventionally do the same. However, baseline glucose for any challenge is not affected by the challenge and could in principle be reported as glucose without specifying the relation to a coming challenge.

We denote the route of the challenge by HL7 Version 2.3 “abbreviations for medication routes” (Table 6). An oral route of administration would be denoted by “PO,”<sup>1</sup> an intravenous route by “IV.”

<b>Abbr.</b>	<b>Challenge Description</b>	<b>Abbr.</b>	<b>Challenge Description</b>
AP	Apply Externally	MM	Mucus Membrane
B	Buccal	NS	Nasal
DT	Dental	NG	Nasogastric
EP	Epidural	NP	Nasal Prongs
ET	Endotracheal Tube	NT	Nasotracheal Tube
GTT	Gastronomy Tube	OP	Ophthalmic
GU	GU Irrigant	OT	Otic
IMR	Immerse (Soak) Body Part	OTH	Other/Miscellaneous
IA	Intra-arterial	PF	Perfusion
IB	Intrabursal	PO	Oral
IC	Intracardiac	PR	Rectal
ICN	Intracervical (uterus)	RM	Rebreather Mask
ID	Intradermal	SD	Soaked Dressing
IH	Inhalation	SC	Subcutaneous
IHA	Intrahepatic Artery	SL	Sublingual
IM	Intramuscular	TRH	Thyrotropin-releasing hormone
IN	Intranasal	TP	Topical
IO	Intraocular	TRA	Tracheostomy
IP	Intraperitoneal	TD	Transdermal
IS	Intrasynovial	TL	Translingual
IT	Intrathecal	UR	Urethral
IU□	Intrauterine	VG	Vaginal
IV	Intravenous	VM	Ventimask

<sup>1</sup> In the United States, PO (an abbreviation for per ora) is used to identify medications taken by mouth.

MTH□	Mouth/Throat	WND	Wound
------	--------------	-----	-------

Examples:

Glucose^pre 100 g glucose PO:MCnc:Pt:Ser/Plas:Qn

Glucose^30M post 100 g glucose PO:MCnc:Pt:Ser/Plas:Qn

Gentamicin^trough:MCnc:Pt:Ser/Plas:Qn

For drug peak (obtained at a time presumed to reflect the highest concentration) and trough (obtained at a time presumed to reflect the lowest concentration) measures the nature of the substance loaded is the same as the analyte name, and need not be included.

2.2.2.2 Physiologic challenges

Some challenges are defined in terms of a physiologic stress, not a dose of a chemical substance. The LOINC names currently cover calorie fasts (no calorie intake), exercise, and fluid restrictions. These challenges are denoted by codes given in Table 7.

In the case of such challenges, the syntax also includes the duration of the challenge.

For example:

post <duration><physiologic challenge>  
Triglyceride^post 12H CFst

Table 7: Example Nature of Challenge	
Type	Description
CFst	Calorie fast. No caloric intake (food) for the period specified in the time part of the term, e.g., POST 12H CFst
Exercise	Exercise undertaken as challenge (can be quantified)
FFst	Fluid "fast." No fluid intake for the period specified

The naming structure is an exact analogous structure to that of chemical challenges. A test for glucose after 12 hours of an energy fast would be represented as:

Glucose^post 12H CFst:MCnc:Pt:Ser/Plas:Qn

A test for osmolality after a 12-hour fluid restriction would be:

Osmolality^post 12H FFst:Osmol:Pt:Urine:Qn

A test for triglyceride after 12-hour energy fast would be:

Triglyceride^post 12H CFst:MCnc:Pt:Ser/Plas:Qn

Two durations can appear in one specification, for example:

Cortisol^1.5H post 0.05-0.15 U insulin/kg IV post 12H CFst:MCnc:Pt:Ser/Plas:Qn

Our rules for naming challenge tests work well only when there is a single intervention followed by a test for one or more components over time. Complex challenge tests involving more than one intervention or complicated sampling techniques need a unique name, but the name may not provide a complete description of all of the test parameters.

### 2.2.2.3 Reporting characteristics of challenge as separate observations

Because we cannot anticipate every type of challenge and route of administration, and because some challenge tests have no usual dose, some challenge tests will not contain a dose. Challenge observations that do not include a specific dose in the name have the word “dose” where a numeric dose would otherwise appear. The general form is:

<analyte>^<time> post dose <route>

Examples:

Glucose^1H post dose insulin IV:MCnc:Pt:Ser/Plas:Qn

The actual dose might then be sent as a comment or as a separate “test” that carries the dose as its value. To accommodate laboratories that wish to transmit the relevant challenge dose as a separate observation, we also define separate test names (and codes) for reporting such doses. This dose could then be sent by the reporting service as a separate result in a separate OBX segment.

The name of the observation that identifies the value of the dose would have the form:

<drug or challenge substance>: <time> post dose <challenge substance>

Examples:

Glucose.PO:Mass:Pt:Dose:Qn

Gentamicin:Mass:Pt:Dose:Qn

Thus we distinguish a drug concentration from the drug dose by means of the system (sample), 4th part, of the test name (see Section 2.5). You can find the observations that carry the dose of drugs or challenges grouped in the class DRUGDOSE in the LOINC database. This approach has the advantages of parsimony and practicality. It also provides an observation ID for the piece of information that must be transmitted along with the request for the observation.

Another example would be:

Oxygen:PPres:Pt:BldA:Qn

Oxygen inhaled:VRat:Pt:Inhl gas:Qn (liters/minute or milliliters/second)

Oxygen inhaled mechanism:Type:Pt:Dose:Nom (to report kind of delivery mechanism, e.g., nasal cannula)

An analogous approach is used for reporting many kinds of associated variables when the variables are not conventionally embedded in the name, in part because there are too many levels of the variables and it is not feasible.

### 2.2.2.4 Generic challenge specifications

We allow for a range of specificity regarding challenges from fully specified to very generic.

Some challenges will be specified fully as described above, e.g., ^30M post 100 g glucose PO . We will

also include: challenges without the amount specified, e.g., ^30M post dose glucose ; those that specify a time elapsed but not a particular challenge, e.g., ^1H post XXX challenge ; those that do not specify the exact time but provide ordering information, e.g., ^2nd specimen post XXX challenge ; or even more generic, ^ post XXX challenge . These latter variants are needed to accommodate challenges that do not fit any common protocol, or referrals to reference laboratories where the study protocol is not reported.

#### 2.2.2.5 Acute and convalescent, pre and post immunization

To assess the efficacy of immunizations, we measure antibody levels before and after the immunization; similarly, we obtain evidence for acute infection by assessing acute and convalescent screens. Both of these cases are reported with the 1st specimen, 2nd specimen syntax, for example:

Acute specimen, 1st specimen, pre-immunization specimen:  
Streptococcus pneumoniae Ab.IgG^1st specimen:ACnc:Pt:Ser:Qn

Convalescent specimen, 2nd specimen, post-immunization specimen:  
Streptococcus pneumoniae Ab.IgG^2nd specimen:ACnc:Pt:Ser:Qn

#### 2.2.3 Adjustments/corrections (3rd subpart)

The third subpart of the data element contains calculations that adjust or correct some measured value. We use this subpart to distinguish corrected or adjusted values from the uncorrected measurement, e.g., corrected cell counts from the raw cell counts. Since these attributes are unique to each measurement, they will be short phrases of text rather than a controlled vocabulary to define the content of the third subpart. However when defined, such a test will have a unique LOINC code and the meaning will be fixed by the text in the third part.

Examples:

Calcium.ionized^^adjusted to pH 7.4:SCnc:Pt:Ser/Plas:Qn

Leukocytes^^corrected for nucleated erythrocytes:NCnc:Pt:Bld:Qn

#### 2.2.4 Distinguishing multiple values for any test via the test name (4th subpart)

HL7 messaging allows for multiple results for one observation. Some systems, however cannot distinguish separate answers per observation, so they made the test names like organism 1, organism 2 or substance 1, substance 2 to report multiple organisms or substances identified in samples. We do not encourage this type of reporting because that distinction can more clearly be accomplished by using one test name (e.g., organism identified) and the HL7 sub ID to distinguish the multiple organisms/substances. However, we have created a few terms to accommodate systems that bind the distinction into their test names. The fourth subpart of the component name will allow reporting of repeat observations taken at the same time and/or on the same specimen.

Example:

Bacteria identified^^^2:Prid:Pt:Stool:Nom:Culture

### 2.3 *Kind of Property (also called kind of quantity) (2nd part)*

The second part of the fully specified name distinguishes between different kinds of quantities relating to the same substance, e.g., the mass concentration versus the substance (molar) concentration of sodium in a urine sample, or the absolute eosinophil count versus the percent of the total white count that is made up of eosinophils. The type of property (kind of quantity) is an IUPAC concept described in the Silver Book<sup>21</sup>. We include most of the relevant IUPAC types of property in the LOINC properties table. (See Appendix F for more detailed examples.)

Main property categories

**Mass:** Observations reported with mass (milligrams, grams, etc.) in the numerator of their units of measure have properties that begin with the word mass: mass content, mass concentration, etc.

**Substance:** Observations reported with moles or milliequivalents in the numerator of their units of measure have properties that begin with the word substance.

**Catalytic activity:** Observations that report enzymatic activity have properties that begin with catalytic, e.g., catalytic concentration, catalytic content.

**Arbitrary:** Results that report arbitrary units in the numerator of their units of measure have a property that begins with arbitrary.

**Number:** Counts are associated with properties that begin with **number**, e.g., a white blood cell count reported as number of WBCs divided by volume of blood, would have a property of Number Concentration.

The pharmaceutical industry has the need for laboratory terms that are not specific as to whether the test measures a substance (substance concentration or substance rate) or mass (mass concentration or mass rate). We have created terms with the properties of MSCnc or MSRat to represent these more general test observations. These will only be displayed in RELMA if the user selects one of two new choices (only MS\* prop, all MS\* prop) on the LIMIT SEARCH screen.

**Category subtypes:** Each of the above major property categories has number of derivatives: **concentration, content, ratio, fraction, and rate** (See LOINC properties table).

**Concentrations:** An amount divided by a volume. These have units such as mg/dL, or gm/L.

**Contents:** An amount divided by a mass. These have units such as mg/gm sample or mg/total protein.

**Ratios:** When a result is reported as one measure divided by another taken from the same system, the property is a ratio. The ratio of the mass concentration of substance A divided by the mass concentration of creatinine in a urine sample, for instance, is a mass concentration ratio (MCrto). The numerator and denominator of a ratio must come from the same system. If the measures come from different specimens, e.g., PT patient/PT control or creatinine serum vs. creatinine urine, it is a relative ratio (RelRto). The ratio of times coming from an actual and normal control (as in some coagulation tests) will be relative time (RelTime), a ratio of mass concentrations coming from two different specimens will be relative mass concentrate (RelMCnc), and a ratio of catalytic concentrations from different specimens will have the property of relative catalytic concentrate (RelCCnc).

**Fractions:** Fractions are ratios of a part over a whole: Creatine kinase.MB/Creatine kinase.total, if measured in grams, is a mass fraction. (Fractions are usually reported as percents.)

**Rates:** A rate is a measure per a time period, e.g., mg/day would be a mass rate (MRat). Clearances have the property of volume rate, but “Clearance” will be included in analyte name to clarify meaning, e.g., Sodium renal clearance:VRat:24H:Urine:Qn

Some measures do not fit the above schema. For instance, IUPAC describes an entitic quantity. This refers to measure per entity (e.g., cells, receptors, and molecules). Entitic quantities usually have units that include the name of some entity, e.g., red blood cells (“per 10<sup>6</sup> RBCs”).

One must be careful when mapping measures of constituents of red blood cells to LOINC code because they can be expressed many ways, e.g., as an amount “per mass of hemoglobin”, “per liter of blood” or “per red blood cell”. The first is a mass content, the second a mass concentration, and the last is an entitic mass (mass per entity) — all different properties.

Some tests report the name of an organism (or initially report the presence of any organism, and later identify the particular strain), toxic substance, antibody or antigen, as a test result. Use “Prid” (presence or identity) as the type of property field for results of this sort.

For example:

Bacteria identified:Prid:Pt:Isolate:Nom:Bacterial subtyping

Barbiturates positive:Prid:Pt:Urine:Nom:Confirm

Correct assignment of properties tends to be the most difficult task for new users of LOINC. Appendix F provides more explanation and many detailed examples.

NOTE: For order sets/panels, the property field may be populated by a dash (-).

Table 8: Example LOINC properties			
Enzymatic Activity		Substance Amount (Moles/Milliequivalents)	
CAct	*Catalytic Activity	RelSCnc	*Relative Substance Concentration
CCnc	Catalytic Concentration	Sub	*Substance Amount
CCrto	Catalytic Concentration Ratio	SCnc	*Substance Concentration
CCnt	*Catalytic Content	ScRto	*Substance Ratio
CFr	*Catalytic Fraction	SCnt	*Substance Content
CRat	Catalytic Rate	SFr	*Substance Fraction
RelCCnc	Relative Catalytic Concentration	SRat	*Substance Rate
		ThrSCnc	Threshold Substance Concentration
	<b>Entitic</b>	SCncDiff	Difference in Substance Concentration
EntCat	*Entitic Catalytic Activity	LsCnc	Log substance concentration

EntLen	Entitic Length	<b>Volumes</b>	
EntMass	Entitic Mass	Vol	*Volume
EntNum	*Entitic Number	VCnt	*Volume Content
EntVol	*Entitic Volume	VFr	*Volume Fraction
EntSub	Entitic Substance	VRat	*Volume Rate
<b>Mass</b>		VRatCnt	Volume Rate Content
Mass	Mass	VRatRto	Volume Rate Ratio
MARic	Mass Aeris	VRto	*Volume Ratio
MCnc	*Mass Concentration	RelVol	Relative Volume
MCrto	Mass Concentration Ratio	RelVRat	Relative Volume Rate
MCnt	Mass Content	ArEnrg	Energy/Area
MFr	*Mass Fraction	ArResis	Resistance/Area
MRat	Mass Rate	ArVol	Volume/Area
MRto	Mass Ratio	<b>Time</b>	
RelMCnc	*Relative Mass Concentration	Time	Time
ThrMCnc	*Threshold Mass Concentration	TmStp	Time Stamp—Date and Time
		TRto	Time Ratio
		TQ2	Timing Quantity 2
<b>Counts</b>		RelTime	*Relative Time
Num	*Number	DateRange	Date Range
Naric	Number Aeris (number per area)	<b>Arbitrary Unit Measures</b>	
NCnc	*Number Concentration (count/vol)	ACnc	Arbitrary Concentration
NCnt	Number Content = Count/Mass	ACnt	Arbitrary Content
NFr	*Number Fraction	ThrACnc	Threshold Arbitrary Concentration
NRat	Number=Count/Time	ARat	Arbitrary Rate
NRto	Number Ratio	LaCnc	Log Arbitrary Concentration
LnRto	Log Number Ratio	RelACnc	Relative Arbitrary Concentration
LnCnc	Log Number Concentration		
<b>Other Properties</b>			
Accel	Acceleration	Hx	History
Addr	Address	Len	Length
Anat	Anatomy	LenFr	Length Fraction
Angle	Angle	LenRto	Length Ratio
Aper	Appearance	Loc	Location
Arb	*Arbitrary	MOM	Multiple of the median
Area	Area	Morph	Morphology
Bib	Bibliographic Citation	OD	Optical density
Circ	Circumference	Osmol	*Osmolality
CircFr	Circumference Fraction	Pn	Patient number
Class	*Class	Prctl	Percentile
Compli	Compliance	Prid	Presence or Identity
CompliRto	Compliance Ratio	PPres	*Pressure (partial)
Cmplx	Complex	Pres	Pressure
Desc	Description	PresRat	Pressure Rate
Diam	Diameter	PressDiff	Pressure Difference

Doc	Document	PresRto	Pressure Ratio
Dosage	Dosage	Quintile	Quintile
Elpot	Electrical Potential (Voltage)	Ratio	Relative Density
ElpotRat	Voltage Rate (=Amperage)	RelRto	Relative Ratio
EmailAddr	E-mail Address	Resis	Resistance
EngCnt	Energy Content	SatFr	*Saturation Fraction
EngFr	Energy Fraction	Seq	Nucleotide sequence
EngRat	Power = Energy/Time	Shape	Shape
EngRatFr	Energy Ratio Fraction	Susc	Susceptibility
EngRto	Energy Ratio	Temp	*Temperature
Enrg	Energy	Tele	Telephone number
Equ	Equation	Txt	Text
Fcn	Function	Threshold	*Threshold
Find	Finding	Titr	Dilution Factor (Titer)
FldResist	Fluid Resistance	Type	Type
Force	Mechanical Force	Vel	*Velocity
Imp	Impression/interpretation of study	VelRat	Velocity Rate
ID	Identifier	VelRto	*Velocity Ratio
Instrct	Instructions	Visc	Viscosity
InvLen	Inverse Length		

\*Starred items are adopted from the IUPAC Silver Book, non-starred items are extensions.

## 2.4 Time Aspect (*Point or moment in time vs. time interval*) (3rd part)

One can either measure a property at a moment (point) in time or measure it over a time interval and integrate, in the mathematical sense, over time. In the latter case, we aggregate a “series” of physiologic states into a single scalar value that reflects some “average” property measured over the specified time interval. Intervals also have relevance for rate measurements such as excretion (substance rate or mass rate) or clearances (volume rates). The amount over an interval is often expressed as a mass rate (MRat, e.g., g/24h) or a substance rate (SRat, e.g., mol/24h). Interval measurements often apply to urine and stool (e.g., collection over 24 hours and calculation of a concentration, total amount, or clearance). They also apply to clinical measurements such as urine outputs where we have shift totals and 24-hour totals. Event counts on physiologic monitors, such as the number of premature ventricular contractions (PVCs) over 24 hours on a Holter monitor, are also of this type.

The allowed values for non-point time aspect are defined as a syntax exactly like the syntax for the times in challenge tests, e.g., <numeric value><S|M|H|W> The most common one is 24H. Table 9 gives some other examples.

For urine collection, 24H is the “standard” integrated measure and these are almost always reported as mass rates (MRat), substance rates (SRat), or catalytic (CRat) rates. These would contrast with spot or random urine tests that are represented as point (PT) measures in our nomenclature and usually reported as concentrations -- MCnc, CCnc, or SCnc for mass, catalytic, and substance concentrations respectively. However, we can also report the average concentration on a 24-hour specimen – in this case the time aspect value would be 24H but the property would be MCnc/SCnc/CCnc instead of MRat/SRat/CRat.

The designation of 24H collection is maintained for tests that traditionally have reference ranges based on amount of substance of a component cleared or excreted in 24 hours. However, a given specimen could

have a 23-hour collection time and would still be called a 24H study. Depending upon the policies and procedures of the lab, they might extrapolate the reported value to what it would have been if the collection continued for the full 24 hours and report it as moles per day.

We also allow indirect specifications of a time window. *Stdy* identifies the duration of the study (without specifying an exact time); *Enctr* identifies the Encounter (ER visit, hospital stay, etc).

Sample volumes reported for timed measurements are carried in other fields or as separate "test" results in other OBX segments.

<b>Table 9: Example Duration Categories</b>					
<b>Abbr.</b>	<b>Duration Descriptions</b>				
Pt	To identify measures at a point in time. This is a synonym for “spot” or “random” as applied to urine measurements.				
Stdy	Duration of the study				
Enctr	Duration of an encounter (hospital stay, visit).				
Episode	Episode				
Gt 1H	Greater than 1 hour				
Ge 1 Hr	Greater than or equal to 1 hour				
Lt 1H	Less than 1 hour				
Procedure dur	Duration of the procedure (surgery, etc.)				
XXX	Not specified; time will be reported in another part of the electronic message				
* (star)	Life of the “unit”. Used for blood products.				
<b>Abbr.</b>	<b>Description</b>	<b>Abbr.</b>	<b>Description</b>	<b>Abbr.</b>	<b>Description</b>
1M	1 minute	7H	7 hours	2W	2 weeks
5M	5 minutes	8H	8 hours	3W	3 weeks
10M	10 minutes	9H	9 hours	4W	4 weeks
15M	15 minutes	10H	10 hours	1MO	1 month (30 days)
20M	20 minutes	12H	12hours	2MO	2 months
30M	30 minutes	18H	18 hours	3MO	3 months
45M	45 minutes	24H	24 hours		
90M	90 minutes	48H	48 hours		
1H	1 hour	1D	1 day		
2H	2 hours	2D	2 day		
2.5H	2.5 hours	3D	3 day		
3H	3 hours	4D	4 day		
4H	4 hours	5D	5 day		
5H	5 hours	6D	6 day		
6H	6 hours	1W	1 week		

#### 2.4.1 Time Aspect Modifier

The second and optional subpart of the time component allows an indication of some sub-selection or integration of the measures taken over the defined period of time: 8H^max heart rate would be the highest heart rate observed over 8H (Shift). Min, max, first, last, mean are the other possible values for this subpart. When nothing is stored in this subpart, we assume a mean value over the time period in questions. Valid values for this subpart are listed in table below.

<b>Table 10: Time Aspect Modifier Codes</b>	
<b>Time</b>	<b>Description</b>
min	Minimum value over interval
max	Maximum value over interval
first	First value observed during an interval
last	Last value observed during an interval
mean	Mean of all of the values observed on the interval (This is the default selection.)

## 2.5 System (Sample) Type (4th part)

System (sample) type is the fourth part of the fully specified test name. It consists of two subparts; the first part names the system, the optional second part, delimited with a “^”, indicates the super system source of the sample if it is not the patient, e.g., fetus, blood product unit, donor, etc.

We define different tests for the combination of component (analyte) and type of system (sample) that are commonly reported. In practice, laboratories include a relatively small range of sample types in their test names. Chemical tests commonly distinguish between serum, urine, blood, and cerebrospinal fluid. Microbiology cultures tend to distinguish between greater numbers of sources.

The first part of the system field should be coded using the abbreviations listed in Table 11. Since this list was defined for reporting sample type in a field of the HL7/ASTM message that is quite independent of the test/measure name, we do not imply that all such types will find their way into distinct LOINC names. However, when a distinction by type of system is required in the name, it should be represented by one of these codes.

For many chemistry tests we have included in the LOINC database a test name for identifying miscellaneous types of body fluid (Body fld), to provide a way to distinguish tests that are performed on fluid types that are not explicitly represented in the database. We use the code XXX to identify a material that is not specified — it could be solid or fluid, for example.

When should we lump a variety of specimen types under the nonspecific code “Body fld” and when we should give a body material its own unique name for a given component? The decision depends upon the degree to which laboratories have reported the system-component pair as a separate “result” and the degree to which the normal ranges for a given component-system have been standardized. By this rule, we will always define different tests for serum and for urine, when a component can be measured in both. We define sweat sodium as a distinct test because it is a standardized test used to diagnose cystic fibrosis. We did not define duodenal fluid sodium as a separate LOINC code because this measure has not been standardized. This does not mean that the specifics about the system would be ignored. It just means that this information would be recorded in another field of the message (the specimen field of the HL7 OBR segment), not in the name. Generally, we will specify the type of system to distinguish at least among blood, urine, cerebrospinal fluid, pleural fluid, synovial fluid, and peritoneal fluid.

For many types of tests, the distinction between plasma and serum is irrelevant. When testing on serum or plasma is clinically equivalent, the system should be recorded as Ser/Plas. Sometimes the test can only be run on either plasma or serum; the component will then be associated with either Ser or Plas in one observation. If the test can be run on either but the results are different and standardized (a very rare circumstance), two separate tests will be defined in our file, one with a system Plas and one with a system Ser. The current LOINC database includes some Ser tests and some Plas tests that should really be Ser/Plas. As we determine that a Ser or Plas test really should have been designated Ser/Plas, we will change the designation.

If the test is run on a combination of types of system (such as a ratio of substance found in CSF and plasma) the codes are joined with a “+”: Plas+CSF, Ser+CSF, Isolate+Ser, etc.

Details about the exact source and collection method (e.g., blood drawn from the right arm and maintained on ice) are not a proper part of the test name and are reported in other parts of the message, e.g., OBX and OBR of the HL7 message.

Table 11: Example Laboratory System/Sample Types					
Abbr.	Name	Abbr.	Name	Abbr.	Name
Abs	Abscess	Fistula	Fistula	Ser	Serum
Amnio fld	Amniotic fluid	Body fld	Body fluid, unsp	Skin	Skin
Anal	Anus	Food	Food sample	Sputum	Sputum
Asp	Aspirate	Gas	Gas	Sppt	Sputum - tracheal aspirate
Bil fld	Bile fluid	Gast fld	Gastric fluid/contents	Stool	Stool = Fecal
BldA	Blood arterial	Genital	Genital	Sweat	Sweat
BldL	Blood bag	Genital fld	Genital fluid	Synv fld	Synovial fluid (Joint fluid)
BldC	Blood capillary	Genital loc	Genital lochia	Tear	Tears
BldCo	Blood – cord	Genital muc	Genital mucus	Thrt	Throat
BldMV	Blood- Mixed Venous	Hair	Hair	Platelets	Thrombocyte (platelet)
BldP	Blood – peripheral	Inhl gas	Inhaled gas	Tiss	Tissue, unspecified
BldV	Blood venous	Isolate	Isolate	Tlgi	Tissue large intestine
Bld.dot	Blood filter paper	WBC	Leukocytes	Tsmi	Tissue small intestine
Bone	Bone	Line	Line	Trachea	Trachea
Brain	Brain	Liver	Liver	Tube	Tube, unspecified
Bronchial	Bronchial	Lung tiss	Lung tissue	Ulc	Ulcer
Burn	Burn	Bone mar	Marrow (bone)	Urethra	Urethra
Calculus	Calculus (=Stone)	Meconium	Meconium	Urine	Urine
Cnl	Cannula	Milk	Milk	Urine sed	Urine sediment
CTp	Catheter tip	Nail	Nail	Unk sub	Unknown substance
CSF	Cerebral spinal fluid	Nose	Nose (nasal passage)	Vag	Vagina
Cvm	Cervical mucus	Nph	Nasopharynx	Vitr fld	Vitreous Fluid
Cvx	Cervix	Penile vessels	Penile vessels	Vomitus	Vomitus
Col	Colostrum	Penis	Penis	Bld	Whole blood
Cnjt	Conjunctiva	Pericard fld	Pericardial fluid	Water	Water
Crn	Cornea	Periton fld	Peritoneal fluid /ascites	Wound	Wound
Dentin	Dentin	Dial fld prt	Peritoneal dialysis fluid	XXX	To be specified in another part of the message
Dial fld	Dialysis fluid	Placent	Placenta		
Dose	Dose med or substance	Plas	Plasma		
Drain	Drain	Plr fld	Pleural fluid (thoracentesis fld)		
Duod fld	Duodenal fluid	PPP	Platelet poor plasma		
Ear	Ear	PRP	Platelet rich plasma		
Endomet	Endometrium	Pus	Pus		
RBC	Erythrocytes	RBCCo	Red Blood Cells Cord		
Eye	Eye	Saliva	Saliva		
Exhl gas	Exhaled gas (=breath)	Semen	Seminal fluid		
Fibroblasts	Fibroblasts				

These abbreviations are used in the laboratory LOINC codes. Systems in clinical LOINC terms are spelled out in full and should be easily understood.

### 2.5.1 Super system (2nd subpart)

The second subpart of the system identifies a “super-system” when it is not the patient, e.g., a blood product unit (BPU), a bone marrow donor, or a fetus. When the super system is not included in a name, “patient” is the assumed default value. This subpart can take on the values in Table 11. Note: we use the term “fetus” broadly to include embryo, placenta and products of conception.

For instance, an example of representing a coagulation study that uses measures on both patient and a control might be:

Coagulation reptilase induced:Time:Pt:PPP:Qn:Coag

Coagulation reptilase induced:Time:Pt:PPP^control:Qn:Coag

Blood banks often report red blood cell antigens for the patient and for each blood product pack assigned to that patient. So we have:

A Ag:ACnc:Pt:RBC:Ord

A Ag:ACnc:Pt:RBC^BPU:Ord

Note: The inclusion of the super system as part of the system represents a change from versions of LOINC prior to Release 1.0K, May 1998. Earlier versions included this information in the (no longer valued) fourth subpart of the component.

## 2.6 Type of Scale (5th part)

The fifth data part of the test name specifies the scale of the measure, and is a required part. The abbreviation of the type of scale (previously called precision), given in Table 12, should be used in the fully specified name. Note that with the release of Version 1.0K, May 1998, we changed the codes for these from SQ to ORD and from QL to NOM to more accurately identify the meaning.

Scale Type	Abbr.	Description
Quantitative	Qn	The result of the test is a numeric value that relates to a continuous numeric scale. Reported either as an integer, a ratio, a real number, or a range. The test result value may optionally contain a relational operator from the set {<=, <, >, >=}. Valid values for a quantitative test are of the form “7”, “-7”, “7.4”, “-7.4”, “7.8912”, “0.125”, “<10”, “<10.15”, “>12000”, 1-10, 1:256
Ordinal	Ord	Ordered categorical responses, e.g., 1+, 2+, 3+; positive, negative; reactive, indeterminate, nonreactive. (Previously named SQ)
Quantitative or Ordinal	OrdQn	Test can be reported as either Ord or Qn, e.g., an antimicrobial susceptibility that can be reported as resistant, intermediate, susceptible or as the mm diameter of the inhibition zone. (Previously named SQN) We discourage the use of OrdQn in other circumstances.
Nominal	Nom	Nominal or categorical responses that do not have a natural ordering. (e.g., names of bacteria, reported as answers, categories of appearance that do not have a natural ordering, such as, yellow, clear, bloody. (Previously named QL)
Narrative	Nar	Text narrative, such as the description of a microscopic part of a surgical papule test.
“Multi”	Multi	Many separate results structured as one text “glob”, and reported as one observation, with or without imbedded display formatting.
Document	Doc	A document which could be in many formats (XML, narrative, etc.)
Set	Set	Used for clinical attachments

Quantitative (Qn) identifies scales that can be tied to some physical quantity through a linear equation. This means that if we have two reports for the same quantity one with a value of 5 and the other a value

of 10 we know that the two are related in amount through the linear equation  $Y = aX + b$ . When the intercept,  $b$ , is non-zero, we have a difference scale. (Fahrenheit temperature is a difference scale.) When it is zero we have a ratio scale (Kelvin temperature is a ratio scale).<sup>22,23</sup> A Qn value may be reported as a value for a “continuous” scale, as is the case for serum sodium, or it may be reported from a series of discrete values, as is the case for titers, e.g., 1:16, 1:32.

Ordinal (Ord): Some observations have values that are well ordered, e.g., “present, absent”, “1+, 2+, 3+”, or “negative, intermediate, positive”, but the values have no linear relationship to one another. We do not know that positive is two or three times as much as intermediate, we just know that positive is more than intermediate. These kinds of observations have an ordinal scale (Ord). Tests with “yes/no” answers are always ordinal (Ord). Tests reported as negative when less than the detection level but as quantified values otherwise should be regarded as quantitative (Qn).

Quantitative/Ordinal (OrdQn): Rarely, a result can be reported in either an ordinal or quantitative scale. The principal example of this scale is a MIC, which can be reported as either resistant/intermediate/susceptible or by the MIC numeric value. The need for terms with OrdQn as scale was further obviated by clarification from HL7 that results such as "POS" and "NEG" should go in the OBX-8 field for normalcy status. Thus, LOINC codes with scale of Qn can be appropriately used in these cases even if the "values" coming back are coded interpretations of the true numeric result value.

Nominal (Nom): Some observations take on values that have no relative order. Think of the numbers on football jerseys. These simply identify the players, they do not provide quantitative information or rank ordering of the players. We refer to these as nominal (Nom) in scale. Blood culture results provide a good example. Possible values could be Escherichia coli (or a code for E. coli) or Staphylococcus aureus. Other examples are admission diagnoses and discharge diagnoses. Any test or measure that looks broadly at patient or specimen and reports the name of what it finds, is a Nom scale. The values of nominal scaled observations are assumed to be taken from a predefined list of codes or from a restricted vocabulary (e.g. a menu of choices). These observations would typically be sent in an HL7 message OBX segment with a Coded Element (CE) data type (in earlier HL7 versions) or its superseding Coded with No Exceptions (CNE) and Coded With Exceptions (CWE) variants (later HL7 versions). It is important to note that the CE and CWE data types allow values to be set as codes with their print text or just as their print text alone. These data types and the Nom scale would not be used for running narrative.

Narrative (Nar): Some observations are reported as free text narrative. The content is not drawn from a formal vocabulary or code system. A dictated present illness would be an example of a scale of narrative (Nar). Many clinical LOINC codes will come in two versions: one for the nominal (coded) version and one for a narrative (free text) version.

We strongly encourage all reporting to be at the most granular level of detail. That is, if three numbers are reported they would each be reported under a unique LOINC code and transmitted in a separate HL7 OBX segment. Occasionally reporting systems are not able to comply with this dictum. For example some chromatography instruments can identify chemicals from the entire spectrum of known chemicals (CAS identifies more than 10 million distinct chemicals) and we may not have specific LOINC codes for reporting out these details. We have designated the scale of Multi to identify results that include many separately structured results as one text “glob” with or without imbedded (display formatting). Some laboratories report all of the details of many multiple measure tests under such globs with test names that correspond to their order name. We strongly discourage such reporting. It defeats the very purpose of individual codes to tag content.

NOTE: Because the individual components of an Order set/Panel often have different scales, the scale for the order set term may be populated by a dash (-).

## 2.7 Type of Method (6th part)

The method by which the test was performed is the sixth part of the test name. Methods need only be expressed as part of the name when they provide a distinction between tests that measure the same component (analyte) but which have different clinical significance or have a different clinical reference ranges. For instance, whole blood glucose tested with a test strip might be distinguished in the method field.

The list of methods given in Table 13 is not exhaustive; we have included only those methods that are abbreviated in the database or which otherwise require explanation or clarification. Most methods are fully spelled out in the database and should be self-explanatory.

Laboratories do not include the method as part of the name for most common chemical and hematological tests. They often need the freedom to choose the instrument according to time of day, urgency of the request for service, availability of the instruments and so on, even though the instruments may employ different methods. The laboratories then adjust each of the “interchangeable” instruments to produce equivalent results even though the instruments may use different methods. Therefore, we do not want to distinguish too finely on the basis of methods. Though method is rarely significant for many chemical and hematological tests, it is often important to immunochemical/serology testing, because the sensitivity and specificity of some tests varies greatly with the method. For this reason, you will commonly see methods included in microbiology tests and coagulation tests within the LOINC database.

This does not mean that information about the method is irrelevant, but that it is not always a meaningful part of the test name. It is an essential element of the internal quality assurance of laboratories.

Remember that both reference range and method can be sent in other fields of ASTM, HL7, and CEN TC251 result messages.

Table 13: Examples of Method Abbreviations		
Method □	Abbr.	Comment □
Agglutination	Aggl	
Coagulation Assay	Coag	To distinguish coagulation assays based on clotting methods
Complement Fixation	Comp fix	
Computerized Tomography	CT	
Cytology Stain	Cyto stain	The staining method used for pap smears, fine needle aspirates and other cell stains.
DNA Nucleic Acid Probe	Probe	See section 2.7.1 for more information about probes.
Chromogenic/Enzymatic Assay	Chromo	To distinguish coagulation assays based on chromogenic (enzymatic) activity.
Enzyme Immunoassay	EIA	Subsumes variants such as ELISA
Flocculation Assay	Floc	
Hemagglutination Inhibition	HAI	
Hemagglutination	HA	Encompasses direct and indirect
Immune Blot	IB	
Immune Fluorescence	IF	Encompasses DFA, IFA, FA
Latex Agglutination	LA	
Leukocyte Histamine Release	LHR	
Minimum Inhibitory Concentration	MIC	Antibiotic susceptibilities
Minimum Lethal Concentration	MLC	Also called MBC (minimum bactericidal concentration)
Molecular Genetics	Molgen	General class of methods used to detect genetic attributes on a molecular basis including RFL, PCR and other methods.
Neutralization	Neut	

Radioimmunoassay	RIA	
Serum Bacterial Titer	SBT	Determines the serum dilution that is capable of killing microorganisms.
Rapid Plasma Reagin	RPR	Microscopic flocculation test, using cardiolipin-lecithin-cholesterol antigen with carbon particles.
Ultrasound	US	
Vertical Auto Profile	VAP	Developed by Atherotech, Inc.
Visual Count	VC	
Venereal Disease Research Laboratory	VDRL	Microscopic flocculation test

### 2.7.1 DNA/RNA probes/measures

We distinguish three kinds of DNA probe methods:

1. Probe without amplification (Probe)
2. Probe with target amplification (Probe.amp.tar)  
See Table 14A for a list of methods that would be identified as Probe.amp.tar in the method part of the LOINC term.
3. Probe with signal amplification (Probe.amp.sig)  
See Table 14B for a list of methods that would be identified as Probe.amp.sig in the method part of the LOINC term.

<b>Table 14A: Examples of specific methods that would be classed as target amplified DNA/RNA Probe.amp.tar (includes nucleic acid target amplification and probe)</b>		
PCR*	Polymerase Chain Reaction	Applies to: DNA, RNA Roche Molecular Systems (thermal cycler) Requires repeated cycles of heating and cooling-each cycle doubles the target
TMA	Transcription Mediated Amplification	Applies to DNA, RNA Gen-Probe, Inc. (isothermal)
NASBA	Nucleic Acid Sequence Based Analysis	Applies to RNA, DNA Organon-Tenika Corp (isothermal)
SDA	Strand Displacement Amplification	Applies to DNA Becton Dickinson (isothermal)
LAT	Ligation-Activated Transcription	
3SR SR	3 Self-Sustaining Sequence Replication	Applies to RNA, DNA Bartel's Diagnostic (isothermal)
LCR	Ligase Chain Reaction	Also probe amplification category method Abbott Laboratories (thermal cycler)
QBR	Q-Beta Replicase or probe amplification category method	Applies to DNA RNA Gene Track Systems. (isothermal)
<b>Table 14B: Examples of specific methods that would be defined in LOINC as signal amplification methods Probe.amp.sig (includes nucleic acid signal amplification and probe)</b>		
HPA	Hybridization Protection Assay	Applies to RNA Gen-Probe Accuprobe
BdnA	Branched Chain DNA	Applies to DNA, RNA Chiron Corp (isothermal)
-----	Hybrid Capture	

\*The items in the first column of the above table are not meant to be used as methods in LOINC terms.

### 2.7.2 Immunofluorescence (IF)

We do not distinguish among many variants of immunofluorescent tests. DFA, ACIF, are all classed as

immunofluorescence (IF).

### 2.7.3 Immune Stain

We classify peroxidase and all other immune stains of tissue under the method category immune stain.

### 2.7.4 Enzyme Immunoassay (EIA)

We classify many variants of enzymes under EIA, including ELISA, CEIA, etc.

### 2.7.5 Coagulation

We distinguish among three kinds of coagulation method: coagulation (Coag), which measures the coagulation activity, immune (Imm), which measures the amount of the coagulant protein, not its activity, and chromogenic (Chromo), which measures the coagulation factor via enzyme rate (also called enzymatic).

### 2.7.6 Stains

We provide very detailed distinctions among various tissue stains, naming them in full. Stain methods that are modifications of a basic method are named using a <basic>. <modification> syntax, e.g., Methenamine silver stain.Jones

### 2.7.7 Clinical measures

We distinguish reported from estimated and measured values; so reported body weight would be the stated weight from a patient or surrogate. Estimated would be the body weight estimated by an observer, and measured body weight.

### 2.7.8 Imaging studies

We distinguish among the major imaging modalities for most measures derived from such imaging studies (e.g., cardiac outputs from a MUGA scan, angiography, 2D Echo, Doppler, etc.).

## **2.8 Short Convenient Names**

As of the August 2002 release of LOINC we have included a new field in the LOINC database called "SHORTNAME". This field will carry a short, mixed case name for the LOINC concept. We have populated these fields for all laboratory and radiology tests. Our goal was to produce names no longer than 30 characters in order to fit within the space allocated by most laboratory reporting systems. In contrast to the formal LOINC name case is significant in the LOINC short name. When possible, we have used common acronyms and common names rather than the more formal name rules of the full LOINC name. For example, we used the English names of allergens in the short names rather than the

formal Latin species names (in part because they were shorter). The LOINC short names are subject to change and should not be used as identifying keys in any database.

These names have been created via a table driven algorithmic process. We have used all upper case to represent acronyms, and mixed case in organism names as specified in naming conventions (e.g., genus is capitalized, species is not). For virus names we used the acronym assigned by Index Virum where available.

## ***2.9 Long Common Names***

LOINC has received periodic requests from users to produce “pretty” display names that could be used in user interfaces, etc. While systematically created names (like the standard LOINC short names) can be guaranteed to be unique, they are sometimes not the most user-friendly. We have always expected that users would link their own local preferred names to LOINC terms for use in reports and displays. In contrast to systematically-created names, user-friendly names are often ambiguous.

After collecting and reviewing display names from several sources, we decided to create a new algorithmically-generated Long Common Name based on patterns we observed. As of the January 2009 release, we have included a new field in the LOINC database called “LONG\_COMMON\_NAME”. These names have been created by an algorithmic process and are checked for uniqueness. Most abbreviations and acronyms that are used in the LOINC database have been fully spelled out in English. For allergens, the common English names are used instead of the more formal Latin species names. For coagulation, the more commonly used phrases such as “Prothrombin time” have been used.

We started creating long common names first for laboratory terms, but are now producing them for all terms. The text strings for the long common names are subject to change over time as we continue to refine the algorithmic process and collect feedback from users. In particular, many of the long common names for clinical terms have not had as intense focus as the laboratory terms have, so we expect these to be refined over time.

## ***2.10 LOINC term names in HL7 messages***

Messaging standards like HL7 typically use a triplet <identifier code>^<descriptive text>^<coding system> for fields that contain coded entries, such as the OBX-3 Observation Identifier field. Given that LOINC now produces at least 3 names for each term (i.e., the six-part Fully-Specified Name, Short Name, and Long Common Name), users have wondered which LOINC name they should use in the <descriptive text> part of that field (or the equivalent displayName attribute in HL7 V3). In general, we recommend the use of the Long Common Name because they are probably the most understandable to human readers. However, the Long Common Names can be quite long and some systems may not be able to accommodate them. In these cases, we would recommend the use of the Short Name because they would fit within the space allocated by most reporting systems, could potentially work as a column name on a flow sheet, and mostly use common acronyms. Using the Fully-Specified Name (e.g. a colon-separated aggregate of the six part name) is generally not recommended because they are not as human friendly and contain more instances of ‘reserved characters’ like “^” and “&”, which would need to be properly escaped in the message.

Furthermore, we recommend the simultaneous communication of the sender’s local code and local name (in addition to the LOINC code and name) as allowed in the messaging structure to facilitate debugging and detection of mis-mappings.

## 2.11 Classes

We assign each LOINC term to a general category called a Class. These categories are relatively broad and are intended to make it easier to sort and browse the database. They are not intended to be binding definitional characteristics of the term, and we may refine them over time. A more detailed listing of the Classes is presented in Appendix B. Throughout this document many of the naming conventions and approaches are described in reference to a Class of terms. Here we provide a bit of explanation about some of the laboratory term classes.

The class of Microbiology includes all tests used to identify microorganisms and evidence for infection by specific organisms as well as cultures direct microscopic exams that identify organisms or prove evidence for present or past infection with specific organisms. Microbiology includes tests for antibodies, antigens, DNA and RNA. The Serology class does not include measures antibodies or antigens related to microorganisms. Molecular pathology class does not include RNA or DNA based tests for infectious organisms. (They are all included in Microbiology.)

The class Blood bank includes all blood bank testing including ABO-Rh testing. Allergy class includes testing for antibodies to allergens (cat dander, trees, etc). Serology includes rheumatological, and autoantibodies, and antigen measures not covered by these two classes. Hematology/cell counts excludes coagulation studies that are found in a separate class. Measures of complement activity are included within Hematology, not Chemistry. Chemistry does not include challenge tests such as Glucose tolerance, ACTH stimulation, etc; these are in a separate category called Challenge tests.

## 3 Special Cases

### 3.1 Findings viewed as variables or as values

For some complex tests there are two ways to organize the results into a report.

#### 3.1.1 Value

Assume a set “X” is made up of five “results” that can have a scale of (absent present) or (0 1). These results could be reported as:

Finding 1 =	Present	- or -	1
Finding 2 =	Absent	- or -	0
Finding 3 =	Present	- or -	1
Finding 4 =	Absent	- or -	0
Finding 5 =	Absent	- or -	0

Each finding is then considered a binary variable. This is sometimes called a “panel” approach.

#### 3.1.2 Variable (Multiple Choice) Approach

The alternative would be to report this information as a single variable (or multiple-choice question) with many possible values:

Variable X - Finding 1, Finding 3

In this case the findings are the values of a variable called Variable X; only the positive findings are

reported as values. Many laboratory tests, e.g., those that test for HLA antigens, red blood cell antigens, or screens for toxic substances, could in theory be presented either way. The microscopic part of the differential count and urinalysis could also be described either way. History and physical findings and (given a real stretch) even culture results could be structured in the panel or multiple choice/multiple answer format.

A single lab may report red blood cell antigens in either way, as a binary panel or a multiple-choice result, depending upon the purpose of the test. The routine cross and type are reported out in the multiple choice pattern format (only positives from a modest fixed set of tested antigens are reported). But if the tests are being used to prove fatherhood, the results are usually reported as a binary panel.

Blood cultures could in theory be regarded as panels:

<b>Test Name</b>	<b>Value</b>
Escherichia coli	absent
Staphylococcus aureus	present
Diphtheroids	absent
Streptococcus pneumoniae	absent
Pseudomonas aeruginosa	present

Although in practice such tests are almost always reported in the multiple choice/multiple answer format, as follows:

<b>Test Name</b>	<b>Values</b>
Blood culture	P. aeruginosa, S. aureus

We bring up these issues to explain why we use a somewhat different data format for some types of tests, and why we sometimes provide for both reporting methods (e.g., HLA blood cell antigen tests) in the LOINC database. When a binary scale is used, the kind of property will usually be arbitrary concentration (ACnc) and the scale ordinal (Ord). When the multiple-choice multiple-answer approach is used, the scale will be nominal (Nom) and the type of property will be presence or identification (Prid).

### **3.2 Blood bank**

Red cell antigens will be named in accordance with the American Association of Blood Banking (AABB) naming standards.<sup>24</sup> In addition to the antigen or antibody, a modifier would be included in the super-system (the second subfield of the SYSTEM field); to indicate whether testing was performed on the patient, donor, or blood pack. Unless explicitly stated, testing is assumed to have been on a material collected from a patient. Additional information about the person identified in the fourth subpart, such as the donor's name or relationship to patient, should be placed in other OBX segments, or comment segments of the message, and would not be part of the test name.

Blood bank reporting illustrates the need for a method of reporting by panel and by multiple-answer mechanism. The LOINC database provides observation names for both kinds of reporting.

#### **3.2.1 Panel reporting:**

Each reportable antigen must have its own test, so that each element in a full set of binary tests could be reported as (negative, positive) or (0, 1).

The fully specified names of A, AB, B, and O blood types (as observations) would be as follows:

Measure of serum antibody against type A blood of donor:

A Ab:ACnc:Pt:Ser/Plas^donor:Ord

Presence of A antigen on donor's red blood cells:

A Ag:ACnc:Pt:RBC^donor:Ord

Presence of A antigen on the blood cells in a pack of blood given to the patient:

A Ag:ACnc:Pt:RBC^BPU:Ord

### 3.2.2 Multiple answer reporting:

All blood antibodies found (or not found) can also be reported in one result term:

Antigens absent:Prid:Pt:BBL^BPU:Nom  
Antibodies identified:Prid:Pt:Ser/Plas:Nom

The LOINC database provides other “observations” for reporting: the status of each blood pack (e.g., held, given, discarded), and for reporting that information when HIS and medical records systems want it; how much of each type of blood product was given at a moment in time; the type of each pack; any adverse reaction to that pack; and the pack number to accommodate laboratories that send this information as discrete observations.

Blood product disposition:Type:Pt:^BPU:Nom

Blood product type:Type:Pt:^BPU:Nom

## 3.3 *Immunocompetence studies (flow cytometry)*

The CD (Cluster of Differentiation) markers in the LOINC database include all of the single markers and the most commonly reported combinations, e.g., CD11C+CD20C+. Most of these are really measuring the number or percent of cells that bear the specific T-cell marker pattern, in which case they should be specified as a subtype of a lymphocyte, e.g., CELLS.CDx. There are other possibilities, and these cell types can also be named; for instance Blasts.CD2 or Abnormal blood cells.CD5.

Two kinds of measures are of interest.

3.3.1 The “absolute” number of such cells per cubic millimeter is represented as number concentrations, for example:

Cells.CD16C+CD56+:NCnc:Pt:Bld:Qn

3.3.2 Percent of cells containing the named marker per 100 cells of that type is represented as number

fraction, for example:

Cells.CD16C+CD56+/100 cells:NFr:Pt:Bld:Qn

The database also includes fully specified names for all of the commonly reported HLA antigens. These are grouped in the class HLA. Experimental methods can define many subtypes of many antigens, so this list is not exhaustive, and is also likely to expand over time.

Example:

HLA-A1:ACnc:Pt:Bld:Ord

### 3.4 *General approach to microbiology results*

The inherently complex structure of the results of microbiological cultures presents unique challenges for the goal of standardized observation names.

**Result Status** (Preliminary, Final) should not be reported as a separate observation or as part of the name. It should be reported in the Result Status field (OBR-25) of the HL7 OBR segment.

**Specimen Type** (Serum, Blood, Urine, etc.) will be indicated in the HL7 OBR segment with the Specimen Source field (OBR-15), but may also be represented in the name.

**Details of specimen collection** will usually be noted as OBX segments or comment segments that accompany the culture result message. The observation identifier for the OBX segment will have the fully specified name of "SPECIMEN COLLECTION DESCRIPTION:FIND:Pt:\*:NOM" and the Observation Sub-ID field will be used to order or group sets of observations. That is, if the material was collected by swabbing a wound of the right upper arm, multiple OBX segments would be created, each with the name "SPECIMEN COLLECTION DESCRIPTION:FIND:Pt:\*:NOM" and the Observation Results fields of the OBX segments would contain respectively "Swab," "Right," "Arm," and "Wound." (The granularity of the actual terms used in the specimen description is at the discretion of the user. Thus, "Right Arm Wound" as the value of a single OBX segment could be used in place of the three codes described in the previous sentence.)

**Descriptions of measurement and culture growth** will be noted as separate OBX segments that accompany the culture result message. The name of the observation identifier will provide the context of the observation. For instance, the name for a quantitative test of bacteria in a specimen would be:

Colony count:Num:Pt:XXX:Qn:VC

**Descriptions of Gram stain findings** will be noted as OBX segments that accompany the culture result message. The name of the observation identifier will be:

Microscopic observation:Prid:Pt:XXX:Nom:Gram stain

The result values that could be reported with this test (which is a multiple-choice, multiple answer type or observation) might include one or more of the following:

Epithelial cells  
Gram-positive cocci in chains

### Many Gram-negative diplococci

**The organisms identified in a culture** will be sent as result values in OBX segments. A separate table of allowable organism names and/or codes is necessary if these are to be sent as understated results. Euzéby's list of bacterial names approximately 20 other authoritative sources (SNOMED is an appropriate source for these organism concepts) may be used as the standard. While "Throat Culture" is the source of the culture inoculum, it is also a label that indicates what kind of media was inoculated and the other techniques used in the laboratory. So, it is a short hand for a kind of method and such will be recorded as the method part of the name. Thus, "Throat Culture", "Blood Culture", and "Clostridium difficile Culture" all represent labels for how a culture was performed. Examples of names of culture results are:

Bacteria identified:Prid:Pt:Bld:Nom:Culture

Bacteria identified:Prid:Pt:Brn:Nom:Culture

Bacteria identified:Prid:Pt:Stool:Nom:Culture

Names of methods of staining directly on a sample/material (where many descriptive observations are possible):

Microscopic observation:Prid:Pt:XXX:Nom:Gram stain

Microscopic observation:Prid:Pt:XXX:Nom:Dry mount

Microscopic observation:Prid:Pt:XXX:Nom:India ink preparation

Microscopic observation:Prid:Pt:XXX:Nom:Trichrome stain

Microscopic observation:Prid:Pt:XXX:Nom:Giemsa stain

Names for results of staining procedures performed on organisms that are growing in culture will use Isolate as the system/sample type. For example:

Fungus identified:Prid:Pt:Isolate:Nom:Fungal subtyping

Names for organism-specific cultures:

Brucella sp identified:Prid:Pt:Bld:Nom:Organism specific culture

Bordetella pertussis identified:ACnc:Pt:Thrt:Ord:Organism specific culture

Chlamydia sp identified:Prid:Pt:Gen:Nom:Organism specific culture

Legionella sp identified:Prid:Pt:Sputum:Nom:Organism specific culture

Note if a test applies to a specific species of organism, the component should include the genus AND species (at least). If the measure applies to a series of species in the same family the string "sp" must be included. If it applies to as subgroup of the genus, then that subgroup should be named.

Names for method for general class of organism:

Fungus identified:Prid:Pt:Wound:Nom:Culture

Bacteria identified:Prid:Pt:CSF:Nom:Culture

Again, the Result Value of these tests would be either organism names or other statements of culture outcome. The table below contains valid values of the culture result from the HL7 OBX segment:

<b>Table 15: Example Culture Results</b>
No growth
Gram-positive cocci
Small Gram negative rod
<i>Escherichia coli</i>
Normal flora
<i>Candida albicans</i>

Presence or Identity (Prid) as a property should be used when the value of a test can identify one set of alternative infectious agents. If the culture is for herpes virus and the culture can have results of herpes virus 1, herpes virus 2, etc., then Prid is the right property. If the culture is for herpes virus and the answer is positive/negative or yes/no, then the property should be arbitrary concentration (ACnc) and the scale ordinal (Ord).

### 3.5 Antimicrobial susceptibilities

The drug susceptibility tests are grouped together in the LOINC database under the class ABXBACT.

Antimicrobial susceptibility tests are named according to the generic name of the drug tested and the methodology used in testing, with property of susceptibility (Susc), and with scale of quantitative (Qn), ordinal (Ord), or OrdQn. Thus, appropriate names would be:

Ampicillin:Susc:Pt:Isolate:OrdQn:MIC

Ampicillin:Susc:Pt:Isolate:OrdQn:Agar diffusion

Ticarcillin+clavulanate:Susc:Pt:Isolate:Qn:MLC

Table 16 lists methods in drug-susceptibility tests.

<b>Table 16: Drug Susceptibility Methods</b>	
<b>Method</b>	<b>Description</b>
Agar diffusion	Bacterial sensitivity via agar diffusion (Kirby-Bauer)
MIC	Minimum inhibitory concentration
MLC	Minimum lethal concentration
SBT	Serum bactericidal titer
Gradient strip	Susceptible by E-Test or gradient strip method

Methodless codes also exist for each antimicrobial agent.

### 3.6 Cell counts

Quantitative counts of various entities and cells in blood, urine, CSF, and other body fluids may be performed and reported in one of three ways. Cell counts in blood are often reported as absolute counts per unit volume (property number concentration, NCnc), or percents of a general cell type, e.g., percent eosinophils, (property number fraction, NFr). Blood cells are usually reported in such a manner, via either a manual or automated count method. Counts on urine and other body fluids can also be done as

direct counts and reported as NCnc or NFr. However, they are more often reported as the number of entities or cells per microscopic high power or low power field, e.g., 5-10 cells per high power field. These are really numbers per area (property Naric). For example, the number of erythrocytes casts per low power field would be reported as:

Erythrocyte casts:Naric:Pt:Urine sed:Qn:Microscopy.light.LPF

Note that even though the values are reported as a range, the scale is still quantitative (Qn), because the values can be related through a ratio. We use HPF or LPF to identify high power and low power fields respectively. Large entities (such as casts) are usually reported per low power fields, smaller entities per high power fields.

One other way such entities are reported is as a pure ordinal, e.g., none, few, moderate, loaded. These would be specified as arbitrary concentration (ACnc) properties with ordinal scale, for example:

Erythrocytes:ACnc:Pt:Semen:Ord:Microscopy.light

### 3.7 Skin tests

These follow the pattern of a challenge test. For a TB skin test it would be:

Tuberculosis reaction wheal^3D post 25 TU ID:Diam:Pt:Skin:Qn

Where TU means tuberculin units, ID means intradermal, Diam indicates a measure of the diameter of the wheal and so on.

### 3.8 Toxicology – Drug of Abuse Screening and Confirmation

Many kinds of test methods are used in toxicology: Screening tests include HPLC, EIA, TLC, RIA, GC, and GCMS (rarely). Confirmation tests are GCMS, LCMS, GC, and HPLC.

Table 17: Drug of Abuse Methods	
Abbr.	Description
HPLC	high pressure liquid chromatography
TLC	thin layer chromatography
GC	gas chromatography
EIA	enzyme immunoassay
RIA	radioimmunoassay
GCMS	gas chromatography/mass spectrometry
LCMS	liquid chromatography/mass spectrometry

Many laboratories use GCMS to signal that the test is a confirmation of a previous screening test, but other methods are also used to confirm, and a given method can be used to screen or to confirm a test. However, it is important that two different methods be used for screen and for confirm and that they both be applied with techniques appropriate to the mode (screen or confirm). So the LOINC committee has determined it is better to distinguish the screening from the confirming procedure by the use of the words “screen” or “confirm,” in the method part of the name, rather than by naming a specific method. Hence LOINC will distinguish toxicology method by Screen and Confirm but not by particular methods.

Toxicology tests can also be performed on a group of drugs/substances or on individual drugs/metabolites/ substances. We will develop LOINC names and codes for both categories: groups of analytes, e.g., “barbiturates” and individual analytes, e.g., “phenobarbital.”

Group test results are usually reported as ordinal (present /absent) but can also be reported as mass concentrations when the numerator is the total mass of the detectable substances in the group. Group tests at the screening level may also be followed by a confirmation at the group level or by confirms of the individual drug/substance tests at the confirmatory level. Individual drugs/substances may be reported as present/absent (Ord) or as mass (or substance) concentrations (Qn).

When individual drugs/substances are reported ordinally, the reporting threshold (the threshold at which a test level is considered positive) may also be reported as a separate “result.” Thus we have separate LOINC codes to report the cutoff used for defining a positive or negative value.

### 3.8.1 Toxicology drug groups

General principles: for each “group” of drugs (amphetamines, benzodiazepines, opiates, etc.) we will define the following kinds of LOINC observations:

#### 3.8.1.1 Screen for a group of drugs/ toxic substances

“X”: ACnc:Pt:Ord:SYS:Screen for the group as a whole  
(Answer = present/absent)

For example, Amphetamines:ACnc:Pt:Urine:Ord:Screen  
Example answer: “present”

Identify the set of drugs/substances screened for by the group test. The answer will be a list of discrete drug/substance names or codes.

“X” tested for:Prid:Pt:SYS:Nom:Screen  
(Answers = individual drugs that this screening test could detect, from a fixed list)

For example, Amphetamines tested for:Prid:Pt:Urine:Nom:Screen (nominal)  
Example answer = “amphetamine, methamphetamine, dextroamphetamine, levoamphetamine, pseudoephedrine”

#### 3.8.1.2 Identify the drugs substances screened for (and perhaps other information). The answer will be a “glob” of narrative text.

“X” tested for:Prid:Pt:SYS:Nar:Screen  
(Answers = individual drugs that this screening test could detect, as a “blob” of text or canned comment)

For example, Amphetamines tested for:Prid:Pt:Urine:Nar:Screen (narrative)  
Example answer = “The EMIT urine screen for amphetamines detects amphetamine, methamphetamine, dextroamphetamine, levoamphetamine as indications of methamphetamine abuse. It is also reactive with a component present in over-the-counter nasal decongestant inhalers, and a positive result must be confirmed by a quantitative method that rules out the non-abuse situation”

When a screen is reported as negative, confirmatory testing is not performed. When a screening test is reported as positive, the result must be confirmed by an independent testing method.

3.8.1.3 Confirmatory testing for the presence of one or more members of the group represented as a single observation.

“X”:ACnc:Pt:SYS:Ord:Confirm  
(Answers = present/absent)

For example, Amphetamines:ACnc:Pt:Urine:Ord:Confirm  
Example answer: “present”

3.8.1.4 List of the actual drug/substances confirmed.

“X” positive:Prid:Pt:SYS:Nom:Confirm  
(Answers = list of analytes detected)

For example, Amphetamines positive:Prid:Pt:Urine:Nom:Confirm  
Example answer: “dextroamphetamine, methamphetamine”

3.8.1.5 More commonly, confirmatory testing is reported as a set of observations, one to report the presence (or quantitative amount detected) of each analyte in the group.

“X”:ACnc:Pt:SYS:Ord:Confirm  
(Answers = present/absent)  
or  
“X”:MCnc:Pt:SYS:Qn:Confirm  
(Answers = quantitative amount)

For example:

Amphetamine:ACnc:Pt:Urine:Ord:Confirm [present]  
Dextroamphetamine:ACnc:Pt:Urine:Ord:Confirm [present]  
Methamphetamine:ACnc:Pt:Urine:Ord:Confirm [present]  
Levomethamphetamine:ACnc:Pt:Urine:Ord:Confirm [present]

3.8.2 Cutoffs

The cutoff levels for screens and confirms of a given substance or group of substances will usually differ. There are three ways to indicate specific cutoffs in LOINC.

3.8.2.1 We provide separate LOINC terms for reporting the cutoff levels of a number of commonly abused substances and substance groups.

“X” cutoff:MCnc:Pt:Urine:Qn:Screen  
“X” cutoff:MCnc:Pt:Urine:Qn:Confirm

For example, Amphetamines cutoff:MCnc:Pt:Urine:Qn:Screen  
Example answer: “1000 ng/ml”

For example, Methamphetamine cutoff:MCnc:Pt:Urine:Qn:Confirm  
Example answer: “500 ng/ml”

3.8.2.2 Two general cutoff terms, one for screen and one for confirm, can be applied to any substance whether or not a pre-coordinated term exists.

XXX cutoff:MCnc:Pt:SYS:Qn:Screen  
 XXX cutoff:MCnc:Pt:SYS:Qn:Confirm

3.8.2.3 For commonly used cutoffs, such as those mandated by regulatory agencies, we provided precoordinated terms for reporting a “present/absent” result with the cutoff specified in the method field:

“X”:ACnc:Pt:SYS:Ord:Screen>“N”  
 “X”:ACnc:Pt:SYS:Ord:Confirm>“N”

For example, Amphetamines:ACnc:Pt:Urine:Ord:Screen>1000 ng/mL  
 Example answer: “not detected”

3.8.3 Reporting the method used for screen and confirm

We provide terms for reporting the method used for screen and confirm tests:

“X” screen method:Prid:Pt:SYS:Nom:\*  
 “X” confirm method:Prid:Pt:SYS:Nom:\*

These would normally be reported in conjunction with terms reporting levels and possibly cutoffs, as in the following example:

Amphetamines:ACnc:Pt:Urine:Ord:Confirm  
 [Answer = positive]  
 Amphetamines cutoff:MCnc:Pt:Urine:Qn:Screen  
 [Answer = 1000 ng/ml]  
 Amphetamines screen method:Prid:Pt:Urine:Nom:\*  
 [Answer = EIA]  
 Amphetamines positive:Prid:Pt:Urine:Nom:Confirm  
 [Answer = amphetamine, methamphetamine]  
 Amphetamine cutoff:MCnc:Pt:Urine:Qn:Confirm  
 [Answer = 500 ng/ml]  
 Methamphetamine cutoff:MCnc:Pt:Urine:Qn:Confirm  
 [Answer = 500 ng/ml]  
 Amphetamines confirm method:Prid:Pt:Urine:Nom:\*  
 [Answer = GC/MS]

3.8.4 Individual drug/metabolite test results

Individual substances can be reported as screens (ordinal), confirms (ordinal) or confirms (quantitative -- usually mass or substance concentrations).

Group test screens may be confirmed by group confirms (as described above) or by individual confirms (Either ordinal or quantitative--depending upon the laboratory's preference)

3.8.4.1 Individual test screen (ordinal)

Methamphetamine:ACnc:Pt:Urine:Ord:Screen  
 Example answer: “present”

3.8.4.2 Individual test confirm (ordinal)

Methamphetamine:ACnc:Pt:Urine:Ord:Confirm  
 Example answer: "present"

### 3.8.4.3 Individual test confirm (quantitative)

Methamphetamine:MCnc:Pt:Urine:Qn:Confirm  
 Example answer: "250 ng/ml"

Individual tests may also be reported as simple quantitative (without confirm or screen), as is the case for therapeutic drug level monitoring.

### 3.8.4.4 Individual substance measured quantitatively; screen/confirm is not relevant

Digoxin:MCnc:Pt:Ser:Plas:Qn  
 Example answer: "1.2 ng/ml"

### 3.8.5 Naming issues

For confirms, would always be looking for specific analytes. For example, you would never look for tetrahydrocannabinol, but would look for delta-9-tetrahydrocannabinol, 11-hydroxycannabinol, etc.

### 3.8.6 Summary

For each "group" LOINC defines the following set of terms:

"Analyte group":ACnc:Pt:Urine:Ord:Screen  
 "Analyte group":ACnc:Pt:Urine:Ord:Confirm  
 "Analyte group":MCnc:Pt:Urine:Qn:Confirm  
 "Analyte group" tested for:Prid:Pt:Urine:Nom:Screen  
 "Analyte group" tested for:Prid:Pt:Urine:Nar:Screen  
 "Analyte group" positive:Prid:Pt:Urine:Nom:Confirm  
 "Analyte group" screen method:Prid:Pt:Urine:Nom:\*  
 "Analyte group" confirm method:Prid:Pt:Urine:Nom:\*

For each individual analyte LOINC now defines the following set of terms:

Analyte:ACnc:Pt:Urine:Ord:Screen  
 Analyte:ACnc:Pt:Urine:Ord:Confirm  
 Analyte:MCnc:Pt:Urine:Qn:Confirm  
 Analyte:MCnc:Pt:Urine:Qn  
 Analyte cutoff:MCnc:Pt:Urine:Qn:Screen  
 Analyte cutoff:MCnc:Pt:Urine:Qn:Confirm

## 3.9 Molecular Genetics LOINC Naming

### 3.9.1 Introduction

Molecular pathology testing can be used for many purposes. In infectious disease testing to identify

organisms and mutations in organisms; in genetic analysis to identify mutations including substitutions, deletions/ insertions, frame shifts and trinucleotide repeats; to identify specific chromosomal translocation and clonality in leukemia and lymphomas; to identify various tumor associated genes and gene deletions; in paternity testing to determine the probability that a person is the parent of a child; and in forensic testing to determine the probability that a criminal is associated with genetic material he/she left as evidence.<sup>25</sup>

### 3.9.2 Terminology

The main methods used are Southern Blot which applies hybridization to selected DNA “chopped” up by restriction enzymes; Northern Blot which applies hybridization to all cellular RNA (which comes naturally in smaller segments) and Restriction Fragment Length Polymorphism (RFLP). RFLP depends on the Variable Number of Tandem Repeats (VNTR) which are normal, but specific variants of each person’s DNA. Southern Blot may be combined with RFLP to target mutations whose exact gene molecular chemistry is not known. For completeness sake, we mention Western Blot, which applies an analogous blot method to protein analysis.

In situ hybridization is a method that applies probes to intact tissue. The cellular patterns of the homologies can then be read microscopically. There are a variety of methods for detecting such in situ probes. One popular method is Fluorescent In-Situ Hybridization (FISH). This technique is analogous to an immune stain except that the molecular binding is based on DNA/RNA homologous instead of antigen-antibody binding.

DNA chips provide a radical new way to identify DNA and RNA sequences. In the patented AFYMETRIX<sup>®</sup> technique, the nucleoside chains are grown using lithography-like methods. Target DNA is tagged with a detector and “washed” over the chip in steps. The locations of the tags on the chip identify the DNA (RNA) in the sample.

Identity testing is used to identify relationships among people and has special complexity. In paternity testing, it can be helpful to have DNA from the child, the putative father and the mother when possible to distinguish the alleles that come from the father.

Blood is the most common specimen for molecular pathology studies. The DNA comes from the leukocytes, bone marrow, tumors, products of conception and forensic specimens are also important specimens.

Forensic testing has special requirements of stringency and often mixes blood antigen testing with RFLP testing. The results are usually reported as a probability.

Genetic changes that occur during the life of the patient such as tumor mutation are called somatic and those that are inherited are referred to as germ line. The nature of the specimen and the testing usually distinguishes these two, so it is not necessary to include this distinction in the test names.

Alleles refer to different forms of a gene. Alleles are distinguished at the phenotype level. Locus refers to a specific DNA (or RNA) codon or the corresponding amino acid in the protein produced by this codon.

The term mutation is usually applied to a genetic variant that causes a functional change in the gene and results in disease. An allele, the term is usually applied to a genetic variant that does not cause a disease.

The string of DNA that codes for a protein is usually interrupted by DNA segments called introns that do

not contribute to the protein definition. Typically the DNA that defines a protein is interrupted by several introns. The coding sequences of DNA between the exons are called introns. Linked together, the exons provide the instructions for creating the specific protein. Exons may be numbered e.g., exon 1, exon 2, etc. Exon numbers sometimes appear in the names of DNA mutations, but for a number of reasons, identifying codon locations relative to an exon is unreliable and we will try to avoid such nomenclature when possible in LOINC names.

A codon refers to the sequence of three nucleotides that code for one amino acid. Codons are numbered from the first codon participating in the protein (in humans the codon for Methionine) starting with codon number 1.

Defects in genes can be coded in one of three different nomenclatures as described in Table 18.

<b>Designation</b>	<b>Explanation</b>
p	Identify the defect by codon by counting the amino acids in the protein produced by the gene counting the first amino acid.
c	Identify the defect by counting nucleotides from the messenger RNA used to produce the protein with intron excluded. These will produce numbers 3x as large as those in the first method.
g	Identify the defect by counting from the first nucleotide in the DNA as it exists as a gene natively in the chromosome with introns included.

### 3.9.3 General Molecular genetics naming rules

When possible, the LOINC component of a molecular pathologic mutation will be named according to the gene name and information about the particular defect (e.g., deleted alanine from position 47). LOINC will resort to the use of the disease name only when the gene has no name and/or the genetic defect is not yet fully specified. We will always include the genetic disease name in the related name field of the database, when the disease is not part of the component; so that users of the database can easily find the LOINC term by the disease name as well.

We use the nomenclature for human gene mutations proposed by Beaudet<sup>26</sup> in the component (when the mutation name belongs in the test name) or as an answer when it belongs as an answer. This nomenclature system recommends that missense mutations be named using single letter amino acid (p-notated - not nucleotide) abbreviations. A list of single letter amino acid codes is given in Table 19.

<b>Amino Acid</b>	<b>Code</b>	<b>Amino Acid</b>	<b>Code</b>
Alanine	A	Leucine	L
Arginine	R	Lysine	K

Asparagine	N	Methionine	M
Aspartic acid	D	Phenylalanine	F
Cysteine	C	Proline	P
Glutamic acid	E	Serine	S
Glutamine	Q	Threonine	T
Glycine	G	Tryptophan	W
Histidine	H	Tyrosine	Y
Isoleucine	I	Valine	V

The system (specimen) used in the LOINC name for genetic testing will usually be BLD/TISS since the distinction between these two specimens is rarely important to the result of a molecular pathology test. We will split this further to accommodate fetal specimens in a later release.

We did not create separate variables for each kind of molecular genetics method, i.e., we will not make up separate variables for measurements done via Southern Blot, PCR, restriction fragment length polymorphism (RFLP) because different methods are only used when they provide the same answer, and the difference is rarely important. Further, a plethora of method variants exists, and we could never hope to keep up with all of these minor variants. Instead, we will use the generic method of MOLGEN (for molecular genetics method) to indicate that a result of the analysis is based on a molecular genetics method rather than some chemical or antigen method.

### 3.9.4 Infectious Diseases

For most infectious disease reporting, the existing LOINC nomenclature (e.g., detecting a particular species of organism by detecting DNA homology) works fine. The word DNA is included as part of the component name and we distinguish the type of method used for detecting the microorganisms (Probe, Probe.amp.tar, Probe.amp.sig). See the Microbiology section for more information.

### 3.9.5 Genetic Diseases

#### 3.9.5.1 DNA diagnostic assays for the detection of specific disease gene mutations.

In most of these cases we require the gene name, the specification of the nomenclature (p, g, or, c) and the mutation name. A LOINC term that identifies a specific mutation will start with the gene name followed by the specification of the mutation in that mutation using Beaudet's syntax. A dot will separate the gene name and the mutation identifier. In general, the form of the component (first part) of the LOINC name will be:

<gene name> gene.<mutation nomenclature><mutation and its location>

For example, Factor V Leiden mutation would be represented as F5 gene.p.R506Q. Where "F5" identifies the gene, "gene" is a fixed part, "p" identifies the kind of mutation nomenclature (protein) and "R506Q" indicates that the amino acid arginine (R) is replaced by glutamine (Q) (see Table 19) at codon #506.

Some examples of fully specified LOINC names for tests of specific mutation are:

F5 gene.p.R506Q:Arb:Pt:BlD:Tiss:Ord:Molgen

Synonyms = Factor V Leiden, Factor V resistance, APC resistance gene

HFE gene.p.C282Y:Arb:Pt:Bld/Tiss:Ord:MoIgen  
Synonyms = HLA-H gene, hemochromatosis gene

CFTR gene.p.F508 del:Arb:Pt:Bld/Tiss:Ord:MoIgen  
Synonyms = Cystic Fibrosis Transmembrane Regulator

The scale used for LOINC codes of this type is Ord. Test procedures that identify single mutations use two DNA probes: one for the normal locus and the other for the abnormal locus. When only the normal probe reacts, the laboratory reports "no mutation". When both the normal and mutation probes react, the laboratory reports "heterozygous". When only the mutation probe reacts it reports "homozygous". Consequently, such single mutation testing produces one of three ordinal "answers":

- a) no mutation
- b) heterozygous mutation (the mutation found in one gene)
- c) homozygous mutation (the mutation was found in both genes in the gene pair)

Specific testing such as this is only possible when the molecular pathology of the gene is very well known and only one defect is being reported.

### 3.9.5.1 DNA diagnostic assays for the detection of multiple disease gene mutations (alleles).

Multiple testing can be reported in 4 styles: a single observation for each pair, two separate observations, gene mutation analysis and narrative.

- a) A separate observation for each pair of genes

This style of reporting is identical to the style used in 3.9.5.1 with each tested mutation having a separate LOINC code. For example:

HFE gene.p.C282Y:Arb:Pt:Bld/Tiss:Ord:MoIgen

HFE gene.p.H63D:Arb:Pt:Bld/Tiss:Ord:MoIgen

- b) Two separate observations.

One observation reports the kind of mutation (allele) found in the first chromosome and another for reporting the kind of mutation for the paired chromosome. In this case, the identity of the allele is reported in the answer. For example

APOE gene allele 1:Prid:Pt:Bld/Tiss:Nom:MoIgen

Answers = E1, E2, E3, or E4

APOE gene allele 2:Prid:Pt:Bld/Tiss:Nom:MoIgen

Answers = E1, E2, E3, or E4

- c) Gene Mutation Analysis

This is really an extension of the above case. The general name is <genetic disease> mutation analysis:Prid:Pt:Bld/Tiss:Nom:MoIgen. The answers are the names of the genes detected. Examples follow:

CFTR gene mutation analysis:Prid:Pt:Bld/Tiss:Nom:MoIgen  
Synonyms = Cystic fibrosis transmembrane regulator

BRCA1 gene mutation analysis:Prid:PT:Bld/Tiss:Nom:MoIgen  
Synonyms = breast cancer risk gene

Answers for these could be “Identifiable Mutation” “Not Identifiable Mutation”

With this type of reporting, a separate observation is usually required to report what alleles or mutations were tested for, so that the person receiving the report will know how to interpret a negative report. In this style of reporting, we may use the disorder name to identify the domain of interest because it covers more than one mutation. The report provides information about multiple possible mutations.

The general form will be

<allele class or disease name> gene mutations tested for:Prid:Pt:Bld/Tiss:Nom:MoIgen.

For example:

CFTR gene mutations tested for:Prid:Pt:Bld/Tiss:Nom:MoIgen  
The answers could include “Delta F508”, “G542X”, “R553X”, “W1282X”, “N1303K”, etc.

#### d) Narrative report

In this case, the information is provided as a bulk narrative report like a visit note and without computer accessible structure. We discourage the use of this approach because it is not useful for automatic analysis.

### 3.9.6 Trinucleotide repeats

A number of diseases, most of which manifest as neurologic disorders are caused by excessive repeats of specific trinucleotides, and the age of onset of the disease is inversely proportional to the number of excess repeats. Examples of these disorders include:

Fragile X syndrome  
Huntington disease  
Spinocerebellar ataxia (SCA1)

We name the component of these terms by the gene when the gene is well defined or the disease, and the name of the trinucleotide that repeats plus the word “repeats”.

<disease name> <trinucleotide> repeats

For example, Huntington disease would be represented as HD gene.CAG repeats

Examples of some fully specified LOINC names are:

FRAXE gene.CGG repeats:Arb:Pt:Bld/Tiss:Ord:MoIgen  
Synonym = Fragile x syndrome

HD gene.CAG repeats:Arb:Pt:Bld/Tiss:Ord:MoIgen  
Synonym = Huntington Disease, It15, Hd, Huntington Chorea

Spinocerebellar ataxia genes.CAG repeats:Arb:Pt:Bld/Tiss:Ord:MoIgen

DMPK gene.CTG repeats:Arb:Pt:Bld/Tiss:Ord:MoIgen

Synonym = Myotonic Dystrophy

These are usually reported “not expanded”, “indeterminate” or “expanded”, so the scale is Ord.

If the actual number of trinucleotide repeats were reported, the property would be entitic number (EntNum) and the scale would be quantitative (Qn). We are not aware of any labs that currently report the actual number. We will define these quantitative variants when they are requested.

### 3.9.6 Hematopathology gene re-arrangement.

Immunocells have an innate genetic variability due to rearrangement. The unique rearrangement can be used to identify the development of a clone of one cell type as occurs in many lymph cell tumors (e.g., lymphoma). We use the following format to identify clonal excess.

Immunoglobulin heavy chain gene rearrangements:Arb:Pt:Bld/Tiss:Ord:MoIgen

Immunoglobulin kappa light chain gene rearrangements:Arb:Pt:Bld/Tiss:Ord:MoIgen

Immunoglobulin lambda light chain gene rearrangements:Arb:Pt:Bld/Tiss:Ord:MoIgen

TCRB gene rearrangements:Arb:Pt:Bld/Tiss:Ord:MoIgen  
Synonym = T cell receptor beta chain

TCRD gene rearrangements:Arb:Pt:Bld/Tiss:Ord:MoIgen  
Synonym = T cell receptor delta chain

TCRG gene rearrangements:Arb:Pt:Bld/Tiss:Ord:MoIgen  
Synonym = T cell receptor gamma chain

These would be reported as “clonal”, or “not clonal”.

### 3.9.7 Translocations

Tests to detect gene-specific translocation breakpoints (with known “partner” genes) should be designated as follows:

T(<breakpoint gene 1>,<breakpoint gene 2>)(<gene1>,<gene2>)gene translocation

For example:

T(9,22)(ABL1,BCR) gene translocation:Arb:Pt:Bld/Tiss:Ord:MoIgen  
Synonyms = Philadelphia chromosome, BCR1, chronic myeloid leukemia, CML

T(14,18)(IGH,BCL2) gene translocation:Arb:Pt:Bld/Tiss:Ord:MoIgen  
Synonyms = Follicular B cell lymphoma, oncogene B-cell leukemia 2, CLL, chronic lymphatic leukemia, follicular lymphoma

T(15,17)(PML,RARA) gene translocation:Arb:Pt:Bld/Tiss:Ord:MoIgen  
Synonyms = RAR, promyelocytic leukemia, myelogenous, retinoic acid receptor, acute promyelocytic leukemia, APL

These can also be expressed as a fraction of cells that have the rearrangement versus total cells of interest:

Cells.t(9,22)(ABL1,BCR)/Cells.total:NFr:Pt:Bld/Tiss:Qn:MoIgen

If specific partner genes are not known, use:

CCND1 gene rearrangements:Arb:Pt:Bld/Tiss:Ord:MoIgen

Synonyms = Lymphoma 1

BCL2 gene rearrangements:Arb:Pt:Bld/Tiss:Ord: Molgen  
Synonyms = Lymphoma 2

The specificity for “major” or “minor” breakpoints should also be designated:

T(9,22)(ABL1,BCR) gene translocation major break points:Arb:Pt:Bld/Tiss:Ord: Molgen

T(9,22)(ABL1,BCR) gene translocation minor break points:Arb:Pt:Bld/Tiss:Ord: Molgen

### 3.9.8 Identity testing

The identity testers usually look at 4 genetic loci (each locus is polymorphic enough that any one match has a 10% error of being incorrect). The loci are independent so if all 4 probes match (including all exclusions and inclusions) the probability of an erroneously match is .0001 (one out of 10,000). They may use more than four depending upon the degree of confidence required by the circumstances of the testing. The forensic community chooses from a set of about 20 probes.

We propose two styles for reporting identity testing: atomic and pre-coordinated definitions

#### 3.9.8.1 Atomic style

This style uses a series of LOINC names to report the kind of index case, the kind of comparison case, the results of the identity testing, and all of the other separate components of the testing. It includes an observation for reporting the actual probes used, and another observation for reporting the population that the probes assume. The method will be MOLGEN.IDENTITY.TESTING. For example:

DNA probes used:Prid:Pt:Index case^comparison case:Nom: Molgen.identity.testing

Population base:Prid:Pt:Probes:Nom: Molgen.identity.testing

Relationship:Type:Pt:index case:Nom: Molgen.identity.testing  
Answers = child, victim, suspect

Relationship:Type:Pt:^comparison case:Nom: Molgen.identity.testing  
Answers = mother, alleged mother, father, alleged father, evidence  
(external to victim)

Confidence of relationship:likelihood:Pt:Index case^comparison case:QN: Molgen.identity.testing  
(this gives the statistical confidence in the conclusion)

Conclusion:Imp:Pt:index case^comparison case:Nar: Molgen.identity.testing  
(this gives summary statement of the conclusion about identity of relatedness)

#### 3.9.8.2 Pre-coordinated definitions alternative

Some of the above atomic terms (e.g., DNA probes used) could also be reported with the pre-coordinated results.

Relationship:likelihood: child^alleged mother:Qn:Molgen.identity.testing  
Synonyms= maternity testing  
(gives the likelihood that the alleged mother is the mother of the index child)

Relationship:likelihood:child^alleged father:Qn: Molgen.identity.testing  
 Synonyms = paternity testing  
 (gives the likelihood that the alleged father is the father of the index child)

Relationship:likelihood: victim^suspect:Qn: Molgen.identity.testing  
 (gives the likelihood that the either the genetic material on the victim is that of the suspect)

Relationship:likelihood: suspect^victim:Qn: Molgen.identity.testing  
 (gives the likelihood that the genetic material on the suspect is that of the victim)

Identity:likelihood:Evidence^suspect:Qn: Molgen.identity.testing  
 (gives the likelihood that the genetic material on the evidence is that of the suspect)

Identity:likelihood:evidence^victim:Qn: Molgen.identity.testing  
 (gives the likelihood that the genetic material on the evidence is that of the victim)

### 3.9.9 Tumor Relation Tumor Genetics

#### Looking at copy number of N-Myc gene (Growth control gene)

N-Myc gene amplification: EntNum:Pt:Bld/Tiss:Qn:Ord: Molgen

N-Myc gene amplification: ArbEnt:Pt:Bld/Tiss:Ord: Molgen  
 Answers = Non-amplified, indeterminant, amplified

(Comment: these are numbers of excess copies resulting from biologic events, not the true measuring process)

#### Gene loss

p gene loss:Arb:Pt:tumor:Ord: Molgen  
 Answer: gene loss, no gene loss

Compare signal from tumor with normal tissue adjusted for total DNA.

## 3.10 Allergy Testing

The allergy testing industry provides tests for more than 450 different allergens today. Most testing looks for IgE antibodies against these allergens. For some allergens testing for IgG and IgA antibodies are available, as well.

For LOINC terms that represent allergen testing, the component is the allergen name plus the type of the antibody (mostly IgE). Most allergens relate to animals, plants or derivatives of such entities. In the past (prior to LOINC vs. 2.04), we used the common name, rather than the scientific name to identify the allergen. However, this approach led to some duplicate term definitions, because two different companies would name the same allergen differently. It also led to ambiguity because two different species of animal or plant would sometimes have the same common name. As of version 2.04, we corrected these problems. To help reduce the ambiguity we now use the Latin name of the species of the biologic entity that causes the allergy.

Some background: First, most allergens can also be identified with a special 2-5 character code assigned by Pharmacia<sup>27</sup> that most allergy testing companies reference in their catalogue of testing. We used these codes to identify duplicate and ambiguous LOINC allergy test terms. These Pharmacia codes are also included in the related names field of the database. Second, allergen tests are often reported in two

styles: a quantitative raw measure and an ordinal (0-6) severity rank (RAST class). LOINC defines separate terms for each of these reporting styles. For example, the two LOINC codes for reporting IgE antibodies to Japanese Millet are:

```
Echinochloa crus-galli Ab.IgE:ACnc:Pt:Ser:Qn
Echinochloa crus-galli Ab.IgE.RAST class:ACnc:Pt:Ser:Ord
```

The RAST class is a categorization of the raw measurement based on specific allergy criteria. The specific IgE class result values (0, 1, 2, 3, 4, 5, or 6) are an ordered categorical response rather than a continuous numeric scale, therefore “RAST class” terms have an ordinal (ORD) scale.

Laboratories also test mixtures of allergens to produce one result. These will be represented in LOINC as follows:

```
(Acer negundo+Quercus alba+Ulmus americana+Populus deltoides+Carya pecan) Ab.IgE:ACnc:Pt:Ser:Ord:Multidisk
Related name = tx2
```

There may be more than one type of allergen for each plant. For instance, IgE antibodies can develop towards tree pollen and the fruit of the same tree. Similarly, antibodies exist for grain and for grain pollen. In these cases, the LOINC component will contain the word “POLLEN” to distinguish the pollen allergen from the food allergen. For example, the LOINC term for corn (maize) IgE antibody would be:

```
Zea mays Ab.IgE:ACnc:Pt:Ser:Qn:
Related names = f8; cultivated corn; maize

Zea mays pollen Ab.IgE:ACnc:Pt:Ser:Qn
Related names: g202: cultivated corn; maize
```

## 4 Clinical observations and measures

### 4.1 Introduction

For most of the measures we include separate observations for summary data, e.g., shift and 24-hour urine output totals. We also provide varying degrees of pre-coordination for the observation, the body site at which it was obtained, and the method. For example, a cardiac output based on the Fick method is distinguished from a cardiac output computed from 2D cardiac echo data.

Physiologic measures are often monitored continuously over time and the instrument reports summary “statistics” over that reporting period. For vital signs these can include minimum, maximum, and mean value over a time period. For intake and output the total is the summary statistic usually reported. When we address measures taken over time, we usually include 1 hour, 8 hour, 10 hour, 12 hour, and 24 hour intervals to cover the varying lengths of work shifts within and across institutions. The LOINC names of these correspond to the form of a 24-hour urine specimen. The times are recorded in the duration (third part) of the name.

The parts of clinical measurement names are largely the same as for laboratory measures, with some subtle differences that are detailed below.

Parts 2, 3, 5 and 6 (type of property, timing, scale, and method) correspond exactly in meaning between

laboratory and clinical LOINC codes.

System: Part 4, body system, has the same general meaning for clinical and laboratory measures, but whereas in the case of laboratory tests the system usually identifies a fluid and a body compartment by implication (e.g., serum, cerebral spinal fluid), for clinical terms, the system is usually a body part (e.g., chest), organ (e.g., heart), or part of an organ (e.g., heart.ventricle). In some cases the system may be an instrument or device attached to the system (e.g., OB ultrasound imaging device).

Component: In the case of laboratory test observations, the component (part 1) usually identifies some chemical moiety that is distributed in the system (glucose, or HIV antibodies). In the case of clinical terms, the component usually identifies a particular projection of a three or four dimension space to a measure of a particular feature (e.g., QRS interval, systolic) of a time changing measure (ventricle.left.outflow tract). In addition, the component is used to distinguish the various ranges or inflections of a physiologic tracing, or to define precisely the section in three-dimensional space in which an area or range is being measured.

The component includes such things as the special kinds of length (e.g., circumference, diameter, or radius) when length is the property, and the specific level and axis on which a measurement of a body part is taken, e.g., circumference taken at the nipple line. The component should remove all ambiguity as to what projection or axis or specific sub-time frame is being measured. So if one is measuring the diameter of the kidney, the system would have to specify kidney.right (or kidney.left), and the component would identify the axis and level at which the diameter was measured (e.g., cross-sectional at level of pelvis). For a measure of chest circumference the system = chest, the component = circumference at nipple line, and the property = length. Areas, lengths, and volumes of organs all have to be specified enough in the component to distinguish a particular area or length that is being measured. When a measure changes over some cycle (e.g., inspiration, expiration, diastole, and systole), then that should also be specified in the component. (Duration is used to identify the duration of an overall study.)

For most clinical measurements, the component is an attribute of a patient or an organ system within a patient. However, attributes of non-patient systems are also often of interest. For example, we might want to know the class of instrument used to obtain the measurement: i.e., the vendor model number or institutional inventory number of an endoscopy. Such identification numbers have a property of ID. Infection control might want the latter reported in order to track nosocomial infections.

When attributes of an instrument or device are being reported, the system is the name of instrument. The same is true when we report characteristics of tubes used to move fluid in and out of body cavities. For example, we might want to report the size and type of a nasogastric tube.

<b>Table 20: Subjects covered to date in clinical LOINC</b>
Body pressure (systolic, diastolic, and mean)
Body measurements
Body weight (and measures used to estimate ideal body weight)
Cardiac ultrasound
Cardiac output, resistance, stroke work, ejection, fraction, etc.
Circumference of chest, thighs, legs
Critical care measures
Dental
Electrocardiographic measures

Emergency department case reports (CDC DEEDS)
Gastroenterology endoscopy
Heart rate (and character of the pulse wave)
Intake and output
Major headings in operative note
Major headings in discharge summary
Major headings of history and physical
Obstetric ultrasound imaging
Ophthalmology measurements
Pathology protocols
Pulmonary ventilator management
Radiology reports
Respiratory rate
Standardization survey instruments
Urology ultrasound imaging

To accommodate the special dimensions of clinical observations we have introduced new options for the kind of property. The new kinds of property are what you might expect from the new kinds of dimensions being measured (e.g., resistance, voltage, work per beat). However, we have also introduced three important new properties:

- Anat** Anatomic is a special case of Prid that identifies anatomic sites.
- Imp** Impression is a diagnostic statement, always an interpretation or abstraction of some other observation (a series of test results, an image, or a total patient), and almost always generated by a professional. (We could also consider the EKG cart's automated diagnoses as impressions.) Impressions are used in laboratory medicine as well as clinical medicine, so you will see them appearing there as well.
- Find** Finding is an atomic clinical observation, not a summary statement as an impression. Physical, historical, review of systems and other such observations have a property of Finding. These may have a scale of Nom for coded findings, Nar for findings reported in narrative text or Ord for ordinal findings.

In clinical measures, super systems (the second subpart of the system component) may be required. For example, we distinguish head measures of a patient versus a fetus as follows:

Circumference.occipital-frontal:Len:Pt:Head:Qn

Diameter.biparietal:Len:Pt:Head^fetus:Qn

#### ***4.2 Atomic versus molecular (pre-coordinated names)***

With clinical terms we almost always have two ways of reporting. Using the first, we can report an observation by reporting a number of atomic variables which together fully describe the observation. For example, we have the following atomic observations for circumference measures. These variables let us deal with all of the unique kinds of circumferences for which we have not yet defined a pre-coordinated term.

Table 21: Examples of Pre-Coordinated Names	
Code	Description
Circumference:Len:Pt:XXX:Qn	The actual measure of some circumference
Circumference site:Anat:Pt:*.Nom	Identifies the body part measured (specifies the system)
Circumference method:Type:Pt:XXX:Nom:*	Identifies the measuring technique used to obtain the circumference (answers = tape measure, derived, imaging)

We also provide pre-coordinated terms that combine some of the atomic variables into one LOINC code. For example, we have:

8279-2 Circumference.at nipple line:Len:Pt:Chest:Qn

and

8293-3 Circumference^inspiration:Len:Pt:Chest:Qn

which provide more specificity and permit the key components of the measure to be expressed as one variable as is the convention in many clinical systems. We call these pre-coordinated codes “molecular” variables.

Within the LOINC database molecular variables will vary with respect to how many atomic components are aggregated. As is true in some laboratory areas, methods often are not included as part of a name, nor are they always reported. The most common molecular aggregation is between functional measure and a particular site of measurement. (e.g., the many different intravascular sites for blood pressure measurements.) But in some cases the molecular variables represent combinations of specific measures and particular methods (e.g., the cardiac output measures). Please note that most molecular variables could also be accompanied by one or more atomic measures to provide special information about the measure, e.g., special circumstances of the measure, or the vendor model number or institutional inventory number of the measuring instrument.

When we have a variable that really reports what would have been contained in the name in a fully pre-coordinated term, we will place an asterisk in the part that will be reported as a value. For example, a variable that is used to report the anatomic site as an atomic variable, would have an asterisk (\*) in the system part of the name. The variable used to report the method of a particular measure would have an asterisk (\*) in the method part of the name.

### 4.3 Radiology Reports

The creation of LOINC codes for naming radiology reports began with a special subgroup of committee members and a collection of report names from a variety of clinical sites. Radiology LOINC codes were first released in 2000. A bolus of over 2,000 new codes were added in December 2004, and the Radiology section of LOINC continues to be an active area of growth.

LOINC names for radiology reports follow the general pattern of other clinical observations and measures, with some subtle differences noted below. Parts, 2, 3, and 5 (type of property, timing, and scale) correspond exactly in meaning to other clinical and laboratory LOINC codes.

### 4.3.1 Diagnostic Radiology Reports

#### 4.3.1.1 Component

Like other clinical LOINC codes, the component identifies a particular projection of a three dimensional space. The component should remove all ambiguity about what projection is being measured.

##### a) Component/Analyte name

The first subpart of the component field delineates the projections and spatial conditions that are present during image acquisition. The first subpart is named using the syntax:

<descriptor> [<number of views>] [<projection beam orientation>] [<body position>]

The <descriptor> identifies the type of images in the report. For diagnostic x-ray and mammography studies, the <descriptor> is either View or Views. For diagnostic ultrasound, MRI, CT, and tomography studies, the <descriptor> is Multisection. The descriptor is the only required field in the component.

Where it is appropriate, additional words are added to the first subpart of the component to clarify the focus of the exam (e.g., Multisection limited, or Multisection for pyloric stenosis).

The <number of views> is an optional parameter to describe a specific integer number of views in the projection. Many radiology report names do not specify the actual projections taken, but rather only the number of views. Some report names describe the number of views in relative terms like “minimum of 3 views” or “less than 4 views”. Where necessary to specify these relative qualifications, we use the following expressions:

- \* Gt = greater than
- \* Ge = greater than or equal to
- \* Lt = less than
- \* Le = less than or equal to

The <projection beam orientation> is an optional parameter that specifies the orientation(s) of the beam with respect to the patient. Widely used abbreviations with unambiguous meanings are employed where appropriate (e.g., PA, AP, etc). Multiple images with different orientations are combined using “&”.

The <body position> is an optional parameter to remove ambiguity about the subject's body position with respect to gravity. Examples positions include prone, upright, supine, for example:

View PA prone:Find:Pt:Abdomen:Nar:XR

In order to accommodate special groupings of views and challenges, where necessary, we will make an exception to the principle of not using parentheses in the component for radiology studies (see section 2.1.3). For example:

Views (AP^standing) & (lateral^W hyperextension):Find:Pt:Knee:Nar:XR

##### b) Report names for portable studies

In general, we do not make names for reports of portable studies, except when the image produced by a

portable study is different than the normal study. For example, portable chest x-ray studies are typically taken at a shorter distance than those taken in the radiology department, and thus we create separate LOINC codes for them:

Views AP portable:Find:Pt:Chest:Nar:XR

#### c) Eponyms and colloquial expressions

Radiology tests are often commonly referred to by eponyms or colloquial expressions. When they are widely used and understood, these names can represent a concise way to communicate the test(s) being reported. In many cases, these expressions convey meaning that spans multiple parameters or even multiple LOINC axes (e.g., COMPONENT, METHOD, and SYSTEM). LOINC names typically employ these expressions only when their meaning is unambiguous, and confine the use of these expressions within one axis. For example:

View Merchants:Find:Pt:Knee:Nar:XR

#### d) Challenge tests

The second subpart of the component is chemical, physical, and/or functional challenges. The naming convention for chemical challenges (e.g., administration of contrast agents) follows the previously described pattern, including abbreviations for route of administration. For example:

Multisection^W & WO contrast IV:Find:Pt:Kidney.bilateral+Collecting system:Nar:XR.tomo

When describing administration of contrast into specific spaces for which abbreviations do not exist, the space is spelled out in full, and preceded by “intra” or “via” according to these guidelines:

We use “intra: when the contrast injected goes directly into this anatomic space, and this space is what is visualized in the study. For example:

Views^W contrast intra lymphatic:Find:Pt:Lymphatics:Nar:XR.fluor

We use “via” when the contrast injected goes through this device (e.g., catheter) and into the anatomic space being visualized. For example:

Views^W contrast via T-tube:Find:Pt:Biliary ducts+Gallbladder:Nar:XR.fluor

Views^W contrast via colostomy:Find:Pt:Colon:Nar:XR.fluor

Physical challenges that are present during imaging are denoted using a similar pattern:

[<existence>] <challenge>

where existence is denoted W, WO, or W & WO. The existence of W & WO denotes separate views, with and without the challenge. For example:

Views^W & WO weight:Find:Pt:Acromioclavicular joint:Nar:XR

#### 4.3.1.1.1 Ambiguity related to “decubitus” in radiology projections and positions

This section describes several issues surrounding radiology naming conventions involving the term

“decubitus” in abdomen and chest x-ray terms, and to describe an LOINC’s accepted naming conventions. The primary point of confusion concerns an ambiguous naming convention that mixes projection and body position.

#### 4.3.1.1.1.1 Accepted Term Definitions

Excerpts from Merrill’s Atlas of Radiographic Positions and Radiologic Procedures<sup>28</sup>:

##### a) Decubitus

Indicates that the patient is lying down and that the central ray is horizontal and parallel with the floor.

Three decubitus positions are named according to the body surface on which the patient is lying:

##### (i) Lateral decubitus (left or right)

In a lateral decubitus position, the patient is side-lying. The position is named left or right by the side of the patient lying on the table.

If the patient’s back is closest to the IR (image receptor, e.g., unexposed x-ray film), this resulting projection is AP. If the patient’s ventral surface (stomach) is closest to the IR, the resulting projection is PA.

The AP projection in the left lateral decubitus position is the most common (and perhaps implied) decubitus view.

However, it is also possible to do a lateral projection in a right or left lateral decubitus (recumbent) position. (Figure 16-17, Merrill, Vol. 3)

The lateral decubitus position is most often used to demonstrate the presence of air-fluid levels or free air in the chest or abdomen because air rises to the right side and views, are not obliterated by air that may be in the stomach.

##### ii) Dorsal decubitus

In a dorsal decubitus position, the patient is supine. The central ray provides a lateral projection. The position can be named left or right by the side of the patient that is closest to the IR.

This is also called a cross-table lateral view (abdomen).

This type of position is commonly used in lateral x rays of the spine when the patient cannot be moved into a standard lateral position and premature infants that cannot be positioned easily.

##### iii) Ventral decubitus

In a ventral decubitus position, the patient is prone; rarely performed, usually in cases of trauma when the patient cannot be moved. The central ray provides a lateral projection. The position can be named left or right by the side of the patient that is closest to the IR.

##### b) KUB

The Kidneys, Ureters, Bladder (KUB) imaging technique is an Abdomen AP projection, often with the patient in the supine position. The KUB view includes anatomical structures from the diaphragm to the symphysis pubis.

#### 4.3.1.1.1.2 Radiology Naming Conventions

In Radiology, LOINC has typically allowed several levels of granularity to accommodate differences in naming conventions (e.g., specifying laterality or not, explicitly specifying contrast use or not). Different levels of granularity have been observed in this domain as well.

Example local term names:

- Abd R Lat Debub XR
- Abd R Decub Port XR
- Abdomen Debuitus
- Chest Decub XR
- Chest L Decub XR
- Xray Chest Decubitus

4.3.1.1.1.2.1 Decubitus is a body position, not a projection. To add clarity to the names, we will use decubitus only to refer to the lateral decubitus position.

a) When using decubitus to specify body position, we will explicitly say “L-lateral-decubitus” or “R-lateral-decubitus”. Including the word “lateral” adds clarity as to which projection we are talking about, and the dashes “-” help link the words together.

(i) Where the intent is to not name a side, we will use “lateral-decubitus”, rather than the more ambiguous, naked “decubitus”.

4.3.1.1.1.2.2 We will not use the term “dorsal decubitus” to refer to the supine position. Supine will be used as a valid body position where needed.

a) Because it is common and clear, we use “lateral crosstable” to mean a lateral projection (right or left) in the supine position, thus encompassing both a projection and body position.

4.3.1.1.1.2.3 The term “ventral decubitus” will not be used to refer to the prone position. Instead, we use prone as a valid body position where needed.

4.3.1.1.1.2.4 Historically, we created some terms in which there was an implied projection (e.g., AP). Through careful review, we revised or deprecated these ambiguous terms so as to make the particular projection explicit include in the name.

a) When a particular projection is not named, it is implied that any potential projection could be done/reported with this code (e.g., AP, PA, or lateral).

- b) An AP L-lateral-decubitus and AP R-lateral-decubitus are considered distinct “views” in our naming conventions. Thus, use plural “views” and not the singular “view” in such terms.
- c) A naked “lateral” in the component means a lateral projection (in any body position).
- d) Historically, LOINC included some terms with the abbreviation KUB as a named view. Through careful review, we have discontinued its use in favor of simply using the projection (AP) and a specified patient position (e.g., supine or upright) where necessary. This avoids the ambiguity about what KUB means with respect to the patient position.
- e) As in other areas tricky spots of Radiology names, parentheses will clarify which projections are being done in which body positions.

#### 4.3.1.2 Timing

Most radiology reports will have a time aspect of “point in time” (PT). A few reports indicate a specific time window (e.g., timed fluoroscopy imaging), and these are named in the usual manner, e.g., <numeric value><S|M|H|W>. Where qualifiers are needed to indicate a relative time frame, we use the following conventions:

- Gt = greater than
- Ge = greater than or equal to
- Lt = less than
- Le = less than or equal to

For example, Le IH

#### 4.3.1.3 System

For all clinical LOINC terms, the system is spelled out in full and should not be ambiguous. For most radiology reports, the system describes what is being viewed, not only the anatomic area of interest. For example, a common study to identify anterior glenoid pathology is the West Point view x-ray. Because this view demonstrates the entire shoulder, not just the glenoid rim, the system is Shoulder:

View West Point:Find:Pt:Shoulder:Nar:XR

We name systems that encompass multiple organ systems by joining them with a “plus” (+). The individual parameters are arranged in cephalocaudal and/or proximodistal order:

Views:Find:Pt:Spine.cervical+Spine.thoracic+Spine.lumbar:Nar:XR

Views:Find:Pt:Spine.lumbar+Sacrum+Coccyx:Nar:XR

While the system describes what is being viewed, it is not an exhaustive list of all structures in the view. For example, in practice, a standard lateral view x-ray of the radius and ulna shows these bones in their entirety as well as the proximal row of carpal bones and the elbow joint. Yet, the system for this report would simply be Radius+Ulna.

#### 4.3.1.3.1 Vessels

For reports of vascular studies, if the system contains multiple vessels, each vessel is named separately and connected by a plus (+), (e.g., Celiac artery+Superior mesenteric artery+Inferior mesenteric artery). If the vessel(s) being viewed is part of a common root, it is named with the common part first, then a dot (.) separator, and then the division (e.g., Vena cava.inferior). If the vessel(s) are independent branches, then they are named independently and connected by a plus (+), (e.g., Superior mesenteric artery+Inferior mesenteric artery).

For studies that view all the vessels in an area, the SYSTEM is typically named in plural form (e.g., Lower extremity vessels, Lower extremity veins). The rationale for this is that most angiography studies demonstrate some vessel branches, not just a single vessel.

#### 4.3.1.3.2 Brain, head, cerebral, and skull

There is presently much variation in radiology system naming patterns pertaining to the anatomical area of the head. We have modeled our naming patterns largely after prevailing conventions. We generally use a system containing Head for reports of MRA, CTA, CT, and US studies. We use the system of Brain with reports of MRI and nuclear medicine studies, and Skull with plain film study reports. For conventional fluoroscopic angiography reports, we use a system containing Cerebral when not specifying a particular artery.

#### 4.3.1.3.3 Extremities

Test names for studies of the extremities often vary in their terminology. The term “arm” technically means the part of the upper limb from shoulder to elbow, but is also commonly used to refer to the entire upper limb, and the term “leg” technically means the part of the lower limb between the knee and ankle, but is also commonly used to refer to the entire lower limb (Dorland’s Illustrated Medical Dictionary<sup>29</sup>). In lieu of this, we have included “arm” and “leg” as broad synonyms, but do not use them as a system. We use “Upper extremity” and “Lower extremity” to refer to the limbs in their entirety or when the visualized region of the limb is not specified. For more specific regions, we name the system based on the anatomy visualized with that particular method. For example, we name an x-ray of the upper arm as:

Views:Find:Pt:Humerus:Nar:XR

#### 4.3.1.3.4 Laterality

For most bilaterally symmetric entities, we create separate LOINC codes for radiology reports differentiated by laterality. Thus, for many studies we have LOINC codes that differ only by the laterality of the system (e.g., Shoulder, Shoulder.left, Shoulder.right, Shoulder.unilateral, and Shoulder.bilateral).

#### 4.3.1.3.5 Series projections with multiple systems

For radiology reports on a series of projections that include multiple systems (e.g., Ribs+Chest), the order the projections are listed in the COMPONENT corresponds with the order of the anatomical sites in the system. In addition, the secondary anatomical site is added to the COMPONENT to clarify which views

were for which anatomical region. For example:

Views lateral & PA chest:Find:Pt:Ribs+Chest:Nar:XR

#### 4.3.1.3.6 Use of dot (.) in system

Using a dot (.) in the SYSTEM signifies that the modifier is a subdivision or component of the main word. No dot (.) is used when the modifier is just an adjective used for clarification. So, we have: Chest.pleura, but Superficial tissue.

#### 4.3.1.3.7 Method

In general, the method for radiology reports corresponds to the method for other LOINC terms. The pattern for naming a radiology method is:

<modality>.[submodality]

##### 4.3.1.3.7.1 Method for angiography terms

LOINC terms use the methods of XR.fluor.angio, MRI.angio, and CT.angio to describe angiography study reports. Radiology systems often use the abbreviations MRA, MRV, CTA, and CTV in test names of angiography studies. Because MRA and CTA can refer to studies of arteries, veins, or both, they are equivalent synonyms to the LOINC methods MRI.angio and CT.angio and are included in the database as synonyms. MRV and CTV are added as synonyms only to terms where the method is MRI.angio or CT.angio and the system contains the word “Vein” or “Veins.”

### 4.3.2 Interventional Radiology Reports

#### 4.3.2.1 Component

Radiology reports for interventional studies under imaging guidance typically contain a component of the form: Guidance for <indication>, where <indication> is description of the nature of the guidance. For example:

Guidance for biopsy:Find:Pt:Breast:Nar:Mam

Guidance for drainage:Find:Pt:Kidney:Nar:US

#### 4.3.2.2 System

The system for interventional radiology reports is named for the anatomical structures being viewed, similar to the pattern for systems of diagnostic radiology reports.

#### 4.3.2.3 Method

In general, the method for interventional radiology reports corresponds to the method for diagnostic radiology and other LOINC terms. The pattern for naming the method is:

<modality>.[submodality]

## 5 Tumor registry

In collaboration with North American Association of Central Cancer Registries, Inc (NAACCR, Inc), we have developed a set of LOINC codes that can be used to communicate tumor registry variables from clinical institutions to tumor registries and among tumor registries. These LOINC terms map to the content of NAACCR data set, and include variables for such things as the hospital at which the tumor was first diagnosed, the primary anatomic site of the tumor, its size, its degree of spread at the time of diagnoses, and a host of other variables of interest to the tumor registries. The NAACCR data set and other cancer-related demographics are identified by the class TUMRRGT.

The NAACCR standards and an implementation guide for transmitting these LOINC tumor registry variables within HL7 messages are available from the NAACCR website: <http://naaccr.org/>.

## 6 Claims attachments

For more information see HIPAA Attachments display in RELMA, the HIPAA Attachment section in RELMA Users' Manual and the respective Claims Attachment books published by HL7 Attachments SIG.

## 7 HL7 LOINC Document Type Vocabulary Domain

This section describes our approach to creating a set of document type codes. This work has been collaboration between the LOINC committee and the HL7 document ontology task force, with initial contributions from Stan Huff, Pavla Frazier, Bob Dolin, Clem McDonald, and continued refinements from many others.

### *7.1 Use of document type codes in HL7 messages*

In creating and maintaining document type codes it is important to distinguish between the purpose of local document names and the names represented by the document type code. Document type codes are created to provide consistent semantics for the names of documents when they are shared or exchanged between independent facilities or enterprises. The names and codes that are used locally within an enterprise are entirely under the control of the local enterprise, and these names are valuable to the work flow and access of information within the enterprise. It is assumed that the exact local name for the document will be retained in the system that created the document and that the local name can be sent along with the document type code when the document is sent to an external organization. The document type code should only express the meaning in a document name that can be shared between independent

organizations.

For example, it is appropriate to have local document names like “Dr. Smith’s Tuesday Pain Clinic Note” or “Albuquerque VA General Medicine Consult Note” for use within an enterprise. However, some parts of these very specific local names are not meaningful outside of the originating enterprise. Thus, proper document type codes would have names like “Outpatient Pain Clinic Note,” or “General Internal Medicine Consult Note.”

<b>Possible local terms</b>	<b>Document type codes</b>
Dr. Smith’s Tuesday Pain Clinic Note	Outpatient Pain Clinic Note
Albuquerque VA General Medicine Consult Note	General Internal Medicine Consult Note

## **7.2 Relationship with terminologies**

### **LOINC**

HL7 will use LOINC codes for clinical document codes, and will not develop an independent document code system for clinical documents. At its option, HL7 may choose to limit its domain to a subset of LOINC codes. HL7 can incorporate any LOINC document code into the HL7 domain.

The naming rules in this document only apply to “clinical notes.” For purpose of this Users’ Guide, a clinical note is a clinical document (as defined by the HL7 CDA Standard), where clinical professionals and trainees produced the document either spontaneously (e.g., I write my admitting note) or in response to a request for consultation. “Clinical Notes” provides a better description of the process.

“Clinical Notes” are to be distinguished from patient reports such as radiology reports, pathology reports, laboratory reports, cardiac catheterization reports, etc., that are generated in response to an order for a specific procedure. Names for most of these later concepts are accommodated well by the clinical LOINC naming structure, and many such codes already exist within the LOINC database.

### **Relationship with HL7 V2.x values**

The HL7 document type code domain will overlap with similar concepts found in HL7 V2.x (user defined table 0270 Document Types; user defined table 0496 Consent Types). Our approach to manage this overlap is:

- Create a mapping from LOINC codes to HL7 V2.x document codes.
- Continue to develop LOINC codes to meet the needs of the HL7 V3 domain that are not present in the V2.x tables.

### **Relationship to a reference terminology**

As soon as possible, the component terms used in the creation of the names of document type codes will be mapped to either the UMLS Metathesaurus or SNOMED CT. This mapping will help to establish the meaning of the terms and will allow aggregation and classification of document type codes based on

definitions, computable relationships, and subsumption hierarchies that exist in the reference terminology.

### 7.3 Elements of Document Type codes

In the following, synonymy or equivalent terms are designated by parenthesis. Document codes are defined by their component parts. The first list of axis values was published in 2003, and served as the basis for an initial set of LOINC codes.

Through both empiric analysis and expert review, we have continued evaluating and refining this list. The following listing contains the current set of axis values for the elements of document type codes that have been vetted by the LOINC Committee. *We are in the process of carefully harmonizing our existing Document terms with these new values.*

#### Kind of Document

Description: Characterizes the general structure of the document at a macro level. Document types are differentiated based on the need to define distinct document headers.

#### Allowed Values:

##### 1. Note

**Description:** Clinical Note – (also known as “Clinical Document”). Documents generated by clinicians as part of patient care, which includes notes written at the initiative of “individual clinic and consulting clinicians.” It does not include clinical reports such as, radiology, pathology, and cardiac catheter reports that are usually stimulated by a particular order. Clinical documents meet five criteria, as defined in CDA 1.0: wholeness, stewardship, authentication, persistence, and human readability.

##### 2. Working draft of additional values for Kind of Document:

Work is presently underway to more fully define the other potential values for Kind of Document. The following list shows the working draft of these values:

1. Administrative note
  - a. Against medical advice note
  - b. Agreement
  - c. Certificate
  - d. Consent
  - e. Contract
2. Advance directive
  - a. Do not resuscitate
    1. Rescinded do not resuscitate
  - b. Living will
  - c. Rescinded advance directive
3. Diagram
4. Flowsheet
5. Legal
6. Letter
7. Note
  - a. Adverse event note
  - b. Alert
8. Report

**Type of Service**

**Description:** Characterizes the kind of service or activity provided to/for the patient (or other subject of the service) as described in the note. Common subclasses of service would be examinations, evaluations, and management. The notion of time sequence, e.g., at the beginning (admission) at the end (discharge) is subsumed in this axis.

1. Communication
2. Conference
  - a. Case Conference
3. Consultation
  1. Confirmatory Consultation
4. Individual Counseling
5. Group Counseling
6. Daily or End of Shift Signout
7. Diagnostic Study
8. Education
  - a. Discharge Instructions
  - b. Discharge Teaching
  - c. Preoperative Teaching
9. Evaluation and Management
  - a. Annual Evaluation
  - b. Assessment
  - c. Crisis Intervention (Psychosocial Crisis Intervention)
  - d. Disease Staging
  - e. Disability Examination
    1. Social Security Administration Compensation Examination
    2. Compensation and Pension Examination
      1. VA Compensation and Pension Acromegaly
      2. VA Compensation and Pension Aid and Attendance or Housebound Exam
      3. VA Compensation and Pension Arrhythmias
      4. VA Compensation and Pension Arteries Veins and Miscellaneous (Misc)
      5. VA Compensation and Pension Audio
      6. VA Compensation and Pension Bones
      7. VA Compensation and Pension Brain and Spinal Cord
      8. VA Compensation and Pension Chronic Fatigue Syndrome
      9. VA Compensation and Pension Cold Injury Protocol
      10. VA Compensation and Pension Cranial Nerves
      11. VA Compensation and Pension Cushing's Syndrome
      12. VA Compensation and Pension Dental and Oral
      13. VA Compensation and Pension Diabetes Mellitus
      14. VA Compensation and Pension Digestive Conditions
      15. VA Compensation and Pension Ear Disease
      16. VA Compensation and Pension Eating Disorders
      17. VA Compensation and Pension Endocrine Diseases
      18. VA Compensation and Pension Epilepsy and Narcolepsy
      19. VA Compensation and Pension Esophagus and Hiatal Hernia
      20. VA Compensation and Pension Eye
      21. VA Compensation and Pension Feet
      22. VA Compensation and Pension Fibromyalgia
      23. VA Compensation and Pension General Medical
      24. VA Compensation and Pension Genitourinary
      25. VA Compensation and Pension Gulf War Protocol
      26. VA Compensation and Pension Gynecological Conditions and Disorders of the Breast
      27. VA Compensation and Pension Hand Thumb and Fingers
      28. VA Compensation and Pension Heart
      29. VA Compensation and Pension Hemic Disorders
      30. VA Compensation and Pension Human Immunodeficiency Virus (HIV)-Related Illness
      31. VA Compensation and Pension Hypertension
      32. VA Compensation and Pension Infectious Immune and Nutritional Disabilities
      33. VA Compensation and Pension Intestines
      34. VA Compensation and Pension Joints (Shoulder Elbow Wrist Hip Knee Ankle)
      35. VA Compensation and Pension Liver Gall Bladder and Pancreas
      36. VA Compensation and Pension Lymphatic Disorders
      37. VA Compensation and Pension Mental Disorders
      38. VA Compensation and Pension Mouth Lips and Tongue
      39. VA Compensation and Pension Multiple Exam

- 40. VA Compensation and Pension Muscles
- 41. VA Compensation and Pension Neurological Disorders
- 42. VA Compensation and Pension Nose Sinus Larynx and Pharynx
- 43. VA Compensation and Pension Peripheral Nerves
- 44. VA Compensation and Pension Post Traumatic Stress Disorder (PTSD) Initial Evaluation
- 45. VA Compensation and Pension Post Traumatic Stress Disorder (PTSD) Review
- 46. VA Compensation and Pension Prisoner of War Protocol
- 47. VA Compensation and Pension Pulmonary Tuberculosis and Mycobacterial Diseases
- 48. VA Compensation and Pension Rectum and Anus
- 49. VA Compensation and Pension Residuals of Amputations
- 50. VA Compensation and Pension Respiratory Diseases
- 51. VA Compensation and Pension Respiratory - Obstructive Restrictive and Interstitial
- 52. VA Compensation and Pension Scars
- 53. VA Compensation and Pension Sense of Smell and Taste
- 54. VA Compensation and Pension Skin Diseases
- 55. VA Compensation and Pension Spine
- 56. VA Compensation and Pension Stomach Duodenum and Peritoneal Adhesions
- 57. VA Compensation and Pension Thyroid and Parathyroid Diseases
- f. Evaluation and Management of a Specific Problem
  - 1. Evaluation and Management of Anticoagulation
  - 2. Evaluation and Management of Hyperlipidemia
  - 3. Evaluation and Management of Hypertension
  - 4. Evaluation and Management of Smoking Cessation
  - 5. Evaluation and Management of Overweight and Obesity
- g. History and Physical
  - 1. Annual History and Physical
  - 2. Admission History and Physical
  - 3. Comprehensive History and Physical
  - 4. Targeted History and Physical
- h. Initial Evaluation
  - 1. Admission Evaluation
  - 2. Admission History and Physical
- i. Plan
  - 1. Treatment Plan
- j. Risk Assessment and Screening
  - 1. Fall Risk Assessment
- k. Subsequent Evaluation
  - 1. Progress Note
- l. Surgical Operation
  - 1. Postoperative Evaluation and Management
  - 2. Preoperative Evaluation and Management
- m. Summarization
  - 1. Transfer Summarization
  - 2. Summary of Death
  - 3. Discharge Note
  - 4. Discharge Plan
  - 5. Discharge Summary
- n. Transplant Candidate Evaluation
- o. Transplant Donor Evaluation
- p. Well Child Visit
- 10. Medication Management
  - a. Medication Reconciliation
- 11. Outreach
- 12. Pathology Procedure
  - a. Autopsy
- 13. Procedure
- 14. Referral
- 15. Respite
- 16. Supervisory Direction
- 17. Triage

## Setting

**Description:** Setting is a modest extension of CMS's (also known as HCFA) coarse definition of settings, which have well defined meanings. Setting is not equivalent to location, which typically has more locally defined meanings and is reported in other parts of the message. Setting would be limited to one of the following categories (with some future extensions possible).

Most clinical report names would include a setting (at least at the top level) to avoid confusion between important classes of reports. For example, The Admission H&P is usually taken to be the Hospital Admission H&P, but it could be confused with the nursing home H&P if not distinguished by the setting. Setting is not a required component of the name.

1. Ambulance
2. Birthing Center
3. Emergency Department
4. Inpatient Hospital
5. Intensive Care Unit
6. Long Term Care Facility
  1. Custodial Care Facility
  2. Nursing Facility
    1. Skilled Nursing Facility
7. Outpatient
  1. Ambulatory Surgical Center
  2. Office
  3. Outpatient Hospital
  4. Urgent Care Center
8. Patient's Home
9. Rehabilitation Hospital
10. Telehealth
11. Telephone Encounter

### Subject Matter Domain (SMD)

**Description:** Characterizes the subject matter domain of a note.

1. Acupuncture
2. Aerospace Medicine
3. Allergy and Immunology
  - a. Clinical and Laboratory Immunology
4. Anesthesiology
  - a. Pain Medicine
5. Audiology
6. Chiropractic Medicine
7. Critical Care Medicine
8. Dentistry
9. Dermatology
  - a. Clinical and Laboratory Dermatological Immunology
  - b. Dermatopathology
  - c. Pediatric Dermatology
10. Emergency Medicine
  - a. Medical Toxicology
  - b. Pediatric Emergency Medicine
  - c. Sports Medicine
  - d. Undersea and Hyperbaric Medicine
11. Ethics
12. Family Medicine
  - a. Adolescent Medicine
  - b. Geriatric Medicine
  - c. Sports Medicine
13. General Medicine
14. Internal Medicine
  - a. Adolescent Medicine
  - b. Cardiovascular Disease
    1. Clinical Cardiac Electrophysiology
    2. Interventional Cardiology
  - c. Endocrinology
    1. Diabetology
    2. Thyroidology

- d. Gastroenterology
  - 1. Hepatology
- e. Geriatric Medicine
- f. Hematology and Oncology
- g. Infectious Disease
- h. Nephrology
- i. Pulmonary Disease
- j. Rheumatology
- k. Sports Medicine
- 15. Medical Genetics
  - a. Clinical Biochemical Genetics
  - b. Clinical Cytogenetics
  - c. Clinical Genetics
  - d. Clinical Molecular Genetics
  - e. Molecular Genetic Pathology
- 16. Mental Health
  - a. Psychiatry
    - 1. Addiction Psychiatry
    - 2. Child and Adolescent Psychiatry
    - 3. Forensic Psychiatry
    - 4. Geriatric Psychiatry
    - 5. Psychosomatic Medicine
  - b. Psychology
- 16. Multi-specialty Program
- 17. Neurological Surgery
- 18. Neurology
  - a. Clinical Neurophysiology
  - b. Neurology Neurodevelopmental Disabilities
  - c. Neurology with Special Qualifications In Child Neurology
  - d. Pain Medicine
  - e. Vascular Neurology
- 19. Nuclear Medicine
- 20. Nutrition Dietetics
- 21. Obstetrics and Gynecology
  - a. Maternal and Fetal Medicine
  - b. Reproductive Endocrinology
- 22. Occupational Therapy
- 23. Ophthalmology
- 24. Optometry
- 25. Oral Surgery
- 26. Orthopedic Surgery
  - a. Orthopedic Sports Medicine
  - b. Surgery of the Hand
- 27. Orthotics Prosthetics
- 28. Otolaryngology
  - a. Neurotology
  - b. Pediatric Otolaryngology
  - c. Plastic Surgery within the Head and Neck
- 29. Palliative Care
- 30. Pastoral Care
- 31. Pathology
  - a. Anatomic and Clinical Pathology
    - 1. Blood Banking Transfusion
    - 2. Dermatopathology
- 32. Pediatrics
  - a. Adolescent Medicine
  - b. Child and Adolescent Psychiatry
  - c. Hepatology
  - d. Medical Toxicology
  - e. Neonatal Perinatal Medicine
  - f. Pediatric Cardiology
  - g. Pediatric Critical Care Medicine
  - h. Pediatric Dermatology
  - i. Pediatric Endocrinology

- j. Pediatric Emergency Medicine
- k. Pediatric Gastroenterology
- l. Pediatric Hematology-Oncology
- m. Pediatric Infectious Diseases
- n. Pediatric Nephrology
- o. Pediatric Otolaryngology
- p. Pediatric Pulmonology
- q. Pediatric Radiology
- r. Pediatric Rehabilitation Medicine
- s. Pediatric Rheumatology
- t. Pediatric Surgery
- u. Sports Medicine
- 33. Pharmacy
- 34. Physical Medicine and Rehabilitation
  - a. Kinesiotherapy
  - b. Pain Medicine
  - c. Pediatric Rehabilitation Medicine
  - d. Spinal Cord Injury Medicine
  - e. Vocational Rehabilitation
- 35. Physical Therapy
- 36. Plastic Surgery
  - a. Plastic Surgery within the Head and Neck
  - b. Surgery of the Hand
- 37. Podiatry
- 38. Preventive Medicine
  - a. Medical Toxicology
  - b. Occupational Medicine
  - c. Undersea and Hyperbaric Medicine
- 39. Primary Care
- 40. Public Health
- 41. Radiology
  - a. Diagnostic Radiology
  - b. Nuclear Radiology
  - c. Pediatric Radiology
  - d. Radiation Oncology
  - e. Radiological Physics
  - f. Vascular and Interventional Radiology
- 42. Recreational Therapy
- 43. Research
- 44. Respiratory Therapy
- 45. Social Work
- 46. Speech Pathology
- 47. Surgery
  - a. Colon and Rectal Surgery
  - b. Pediatric Surgery
  - c. Surgery of the Hand
  - d. Surgical Critical Care
  - e. Thoracic Surgery
  - f. Transplant Surgery
  - g. Vascular Surgery
- 48. Tumor Board
- 49. Urology

## Role

**Description:** Characterizes the training or professional level of the author of the document, but does not break down to specialty or subspecialty.

- 1. Assistant
- 2. Case Manager
- 3. Clerical
- 4. Counselor
- 5. Fiduciary
- 6. Interdisciplinary
  - a. Team
- 7. Medical Assistant
- 8. Nursing

- a. CRNA
- b. Certified Nursing Assistant
- c. Clinical Nurse Specialist
- d. Nurse Midwife
- e. Nurse Practitioner
- f. Licensed Practical Nurse
- g. Registered Nurse
- 9. Patient
- 10. Physician
  - a. Attending
  - b. Fellow
  - c. Intern
  - d. Resident
- 11. Physician Assistant
- 12. Student
  - a. Sub Intern
- 13. Technician
- 14. Therapist

\* Physician subsumes medical physicians and osteopathic physicians.

### 7.4 Rules for Creating Clinical Notes from Multiple Components

Names for required clinical notes would be constructed by picking entries **from the Kind of Document axis and at least one of the other four axes**. The LOINC committee will create LOINC codes for all required combinations (not all possible combinations).

The original document ontology terms were created only for the document type of “note” and with the general naming pattern:

<Subject Matter Domain> : <Training / Professional Level>: <Setting>: <Type> : Note

As we have revised and refined the elemental axes in the document ontology, simple names would be constructed and ordered as follows:

Table 23. Document Ontology LOINC Naming Rules						
Component	Property	Time	System	Scale	Method	Class
<Type of Service> <Kind of Document>	Find	Pt	<Setting>	Doc	<SMD>.<Role>	DOC.CLINRPT

In general, combinations from within an axis are allowed in a term name where they make sense (SMD, Service), but are disallowed where they do not (Role, Setting). Combinations will be represented with a plus (+), so as to distinguish from elements containing “and” or “&”. Where a particular element is not defined for a given term and leaves a LOINC axis blank, the LOINC name will include the {} naming convention. For example, if a Setting is not designated, the System will be “{Setting}”.

LOINC codes for clinical notes designed according to this model and are assigned a class of DOC.CLINRPT.

Example LOINC codes in the Document Ontology include:

<b>Component</b>	<b>Property</b>	<b>Time</b>	<b>System</b>	<b>Scale</b>	<b>Method</b>	<b>Class</b>
Group counseling note	Find	Pt	Hospital	Doc	{Provider}	DOC.CLINRPT
Evaluation and management note	Find	Pt	Outpatient	Doc	{Provider}	DOC.CLINRPT
Evaluation and management note	Find	Pt	{Setting}	Doc	{Provider}	DOC.CLINRPT
History and physical note	Find	Pt	{Setting}	Doc	{Provider}	DOC.CLINRPT

## 7.5 Future Work

We continue active development and refinement of the Kind of Document axis. As we continue this work, we intend to develop equally specific definitive documents for other kinds of health case associated documents.

## 8 Order Panels (Batteries)

Beginning with version 1.0o, the LOINC database was expanded to include order sets/panels. These have been identified with the word "PANEL" in the component name. Since the property type will vary depending on the panel elements, the second part of the LOINC name may be populated by a dash (-). The scale (5th part of the LOINC name) will be populated by a dash (-) if the panel elements could have different scales.

If a government authority recognizes the order set, it will include the year that an order set took effect. For example:

Comprehensive metabolic 2000 panel.

Using RELMA, you can view the list of the individual test components included in each panel (order set). The elements will be accompanied by a flag that will denote the expected appearance of the panel element in the panel when resulted. A flag is always one of three states:

- R - Required. The panel element is always expected to be reported when the panel is resulted.
- O - Optional. The panel element may not be reported with a panel result depending upon institutional policy or capabilities of the reporting lab.
- C - Conditional. The panel element is a key finding in the panel report and should be assumed to be negative, absent or not present if the panel result does not include data for this element.

Some example order sets:

<b>24358-4</b>	<b>Hemogram WO platelets panel</b>	<b>-</b>	<b>Pt</b>	<b>Bld</b>	<b>Qn</b>	<b>R</b>
26464-8	Leukocytes	NCnc	Pt	Bld	Qn	R
26453-1	Erythrocytes	NCnc	Pt	Bld	Qn	R
718-7	Hemoglobin	MCnc	Pt	Bld	Qn	R
20570-8	Hematocrit	VFr	Pt	Bld	Qn	R

30428-7	Mean corpuscular volume	EntVol	Pt	RBC	Qn	R
28539-5	Erythrocyte mean corpuscular hemoglobin	EntMass	Pt	RBC	Qn	R
28540-3	Erythrocyte mean hemoglobin concentration	MCnc	Pt	RBC	Qn	R
30384-2	Erythrocyte distribution width	EntVol	Pt	RBC	Qn	O
30385-9	Erythrocyte distribution width	Ratio	Pt	RBC	Qn	O
<b>24326-1</b>	<b>Electrolytes 1998 panel</b>	-	<b>Pt</b>	<b>Ser/Plas</b>	<b>Qn</b>	<b>R</b>
2951-2	Sodium	SCnc	Pt	Ser/Plas	Qn	R
2823-3	Potassium	SCnc	Pt	Ser/Plas	Qn	R
2075-0	Chloride	SCnc	Pt	Ser/Plas	Qn	R
2028-9	Carbon dioxide	SCnc	Pt	Ser/Plas	Qn	R
330373	Anion gap	SCnc	Pt	Ser/Plas	Qn	O

LOINC has created a series of spreadsheets that contains sets of related panels. These spreadsheets may include information on multiple panels or just a single panel. Each spreadsheet contains three worksheets: list of LOINC terms within the union of the panels in the package, a table that defines the nesting of the panels within a given panel, and a table that defines all of the answer lists in the panel. This content can be easily used to load a database, but every panel in LOINC is not yet represented this way. You can also get to the panels content by going to the panel screen in RELMA. You can look at the content of every panel on this screen in report format. You can also pick any panel in the grid of a search, right click, and then export.

## 8.1 Goals

We have gotten many requests for a standard set of test order codes from Medical Information System vendors. They want standard codes for the common orders so they can install their system with a set of usable starter set of order codes. They also want them to ease the cross communications among merging hospitals.

LOINC codes have been defined for most individual laboratory observations and for many clinical observations, and claims attachments. Obviously, these same LOINC codes can be used to order individual laboratory and clinical observations, as well as to report the LOINC code for Blood Hemoglobin (LOINC # 718-7) could as easily be used to order a Blood Hemoglobin, as well as to report the result of that test. Pre-existing LOINC codes could also be used to order more complex observations. The Urinary Creatinine Clearance (LOINC # 2164-2) could also be used order code Creatinine Clearance. Since the calculation of creatinine clearance requires two distinct measures (serum creatinine and 24-hour urine creatinine), an order for creatinine clearance implies an order for these two other measures. However, the existing single value LOINC codes could not be used to order many laboratory and clinical procedures that are ordered as a single-named test (battery), such as CBC, urine dipsticks, blood differential count, LDH isoenzymes. Similarly physicians order Blood pressure measures and expect to get (at least) the diastolic blood pressure and the systolic blood pressure. Though these are separate observations, for practical purposes one is never measured without the other. Initially, we created LOINC codes for the common “fixed” observation packages. By fixed, we mean that certain kinds of measures will always be part of the battery, and the production of that particular set of measurements is tightly bound to the procedure or instruments that produce the values and or by a government mandate (e.g., LOINC # 24325-3: Hepatic function HCFA 2000 panel). Other types of order codes have evolved.

**Background on kinds of results found in order sets**

To understand the rules about creating order sets, we distinguish several kinds of results in orderable test batteries (or sets).

**8.2 Reflex tests**

Testing can be done in steps. A certain number of analyses are done at the first step, then depending upon the values of those analytes different analyses (observations) are performed. For example, a TSH test might be done first and depending upon its value, other confirmatory tests would be done. We have not yet addressed the naming of Panels with reflex components in LOINC. This is work for the future.

**8.3 Calculated or derived results**

The results in an order set often include results that simple calculations based on the primary measurements. For example, it might include the absolute concentration and the percent concentration of a given element, such as basophils. In the information theoretic sense these do not provide additional information. So we will usually use one order panel name regardless of how many values were calculated from the primary measurement.

**8.4 Associated observations**

Some sets consist of a set of measures produced by the laboratory and a set of observations obtained by the placer and sent along with the request. For example, placers will usually report the percent inspired O<sub>2</sub> when they request an arterial blood gas and the laboratory reports that value along with the values it measures directly. We call these “associated observations” and count the volumes and times of collection in this category for the purpose of this discussion. We will not define distinct order panels that vary with the number of clinical variables (not measured by the lab) that are included in the report.

**8.5 LOINC Rules for representing order panel names**

We will use most of the same general LOINC naming rules for Batteries of Observations (Panels) as for individual observations.

Component Name: For orders sets consisting of three or more constituent tests, the component name will be a concatenation of:

- (1) A name (e.g., Hemogram, Differential count, Vital Signs) to convey the content of the panel
- (2) The word “Panel” included to unambiguously identify that this LOINC term refers to a panel or battery

In the case that a well-defined panel exists but has no conventional name, we will include each of the distinct measured entities separated by ampersand (&) in the component name. So for example, when a creatinine is measured along with sodium in a 24-hour urine, we will use this convention to build up panels from other panels. We may also use a more efficiently syntax, which implies repeat of the first part of the name, e.g., Chlamydia Ab IgM & IgG Panel.

Any of these batteries may variously include in the report a variety of other values derived from the reported measures, information sent along with the request (e.g., inspired O<sub>2</sub> for blood gases). In most

cases we will not make up different names for the same set of tests done by different methods. Because of the possible mixtures of methods within a panel, representing these distinctions would cause an explosion of the distinct Panel, which would (usually) be a burden on the ordering provider. Further, in a given setting the ordering provider can only order the methods that are provided by his usual producer. Implied in the order is “Give me the battery produced by your usual methods”. In special circumstances, we might provide method specific observation panels, e.g., when blood pressure is usually done by automated methods, the provider might want the option of obtaining a blood pressure by manual methods as a double check.

Property, Timing, Scale and Method: We will not usually value the property type (the second part of a LOINC name) of an order panel because the property varies within the measures included in a battery. But since this field cannot be null in a LOINC name, we will include a dash (-) in this field, but we will usually value the timing and the system and the scale field.

At this first phase we have defined batteries for:

- Hemograms and differential counts (both automated and manual)
- Arterial blood gases
- Urinalyses
- Isoenzymes
- Antibodies for IgG and IgM when they are done in pairs
- Common toxicology batteries
- Susceptibility testing
- Chemical batteries defined by HCFA
- A few clinical orders

Description of some LOINC Panels (Order Set Names):

LOINC_NUM	LOINC Fully Specified Name	Description
24358-4	Hemogram WO platelets panel:-:Pt:Bld:Qn	HCT & HGB & WBC & RBC & Indices
24359-2	Hemogram WO platelets & W manual differential panel :-:Pt:Bld:Qn	Hemogram & Differential Count
24317-0	Hemogram & platelets WO differential panel:-:Pt:Bld:Qn	HCT & HGB & WBC & RBC & Indices & Platelets
24338-6	Gas panel:-:Pt:Bld:Qn	pH & PO <sub>2</sub> & PCO <sub>2</sub> on blood without specifying whether arterial, venous, or other source. The report would usually include an observation about the inspired O <sub>2</sub> sent along with the report. It may include a variety of other patient characteristics sent by the requester and a variety of computed variables.
24336-0	Gas panel:-:Pt:BldA: Qn	pH & PO <sub>2</sub> & PCO <sub>2</sub> on arterial blood. The report would usually include an observation about the inspired O <sub>2</sub> sent along with the report. It may include a variety of other patient characteristics sent by the requester and a variety of computed variables.
24339-4	Gas panel:-:Pt:BldV:Qn	pH & PO <sub>2</sub> & PCO <sub>2</sub> on venous blood. The report would usually include an observation about the inspired O <sub>2</sub> sent along with the report. It may include a variety of other patient characteristics sent by the requester and a variety of computed variables.
29274-8	Vital signs measurement:Find:Pt:^Patient^Multi	Diastolic Blood Pressure & Systolic Blood Pressure & Pulse Rate & Respiratory Rate
24357-6	Urinalysis macro (dipstick) panel:-:Pt:Urine:-	Urinalysis dip stick results. Usually includes Glucose, Bilirubin, estimate of leukocytes, estimate of RBCs, estimate of bacteria, Ph, Specific gravity. But we do not make distinctions about the exact set of measures on the dipstick. The ordering clinician will not necessarily know what particular dipstick is being used

		and is not able or interested in making those distinctions.
29576-6	Bacterial susceptibility panel:-:Pt:Isolate:OrdQn	Would include susceptibility results for the antibiotics relevant to the isolates and the kind of culture.

## 9 Evolving principles for naming collections

### 9.1 Goals and general approach

We are in the process of evolving our model for naming collections in LOINC. Our goals in refining this model are to:

- Create names that are consistent across different subject domains within LOINC
- Make it easy to create a list of all codes that could be used as document type in CDA
- Make it easy to create a list of all codes that could be used as section headings in CDA
- Avoid proliferating names

To this end, we are developing rules for naming of collections will apply to both laboratory collections (CBC, CHEM7) coded and structured clinical collections (Vital Signs), documents (Admit History and Physical Exam), Apgar scores, Braden Scale, Pain scales, etc. There will be two categories of names for collections:

- Names for panels with enumerated discrete contents, and
- Names for general collections of information.

Using the existing panel mechanisms, the LOINC database will record the association between LOINC collections and individual observations where these associations are known. For example, LOINC already records the expected contents for CBC, Liver Enzymes, etc. It will also include definitions for Vital Signs, Cardiac Catheterization, Braden Scale, surveys, etc. We will create a single LOINC code for any general collection of information where the information content of the collection is the same, regardless of whether the content is a text document, a scanned image of text, or a sound file of the same information.

Since collections are named by their real or anticipated contents, the same LOINC code could be used as either a document type or as a section type.

### 9.2 Collections as orders and observations

The same LOINC code will be used for ordering a procedure, naming the document produced as the description of the procedure, or naming the structured and coded set of observations from the procedure.

For panels, the same code for CBC would be used as the ordered item in an order record or message, and as the panel identifier in the OBR segment of a result record or message. The same pattern would be followed for laboratory procedures and clinical procedures.

For general collections, the same code would be used as the ordered item in an order record or message, and as the result identifier in a result message. For example, the general collection name could be used in a result message as the identifier of a document type, as a section label, as the universal identifier in an OBR segment, or as the identifier in an OBX segment depending on the circumstances. The same pattern

would be followed for radiology procedures and clinical procedures.

We are not taking away the flexibility of having the ordered code be different from the result code. For example, it is often desirable for the order code to be less specific and more abstract than the result code. LOINC would contain codes for something like “Exercise EKG” with the expectation that the result could come back as “AHA Protocol Stress EKG Result”. The point is that when appropriate we would use the same LOINC code in the contexts of orders and results. We would NOT make LOINC codes that meant “CBC Order” and “CBC Result”, we would use the same LOINC code for CBC in both orders and results.

Current practice would also continue where a “pure” procedure is ordered and discreet results would be returned. For example, Urine Microscopic Exam could be ordered and discreet values for cell types, casts, amorphous material, etc. would be returned.

### 9.3 LOINC SCALE for collections

The SCALE for panels will be “Panel.” The SCALE for general information collections will be “Doc”, short for document, which is used in the most general sense of a text document, image, scanned text image, etc. “Doc” would replace the current use of Nar (narrative) or Nom (nominal) for general information collections in the current LOINC database.

The LOINC committee will review current contents of the LOINC database and modify names appropriately to conform to the new conventions. We will not implement the name changes until after the current Attachments NPRM is final.

### 9.4 Examples of proposed changes according to new policy

Table 27: Example of Proposed Changes							
LOINC#	Component	Property	Time	System	Scale	Method	Class/Type
24358-4	Hemogram panel	-	Pt	Bld	Panel		PANEL.HEM/BC
24320-4	Basic Metabolic HCFA 98 panel	-	Pt	Ser/Plas	Panel		PANEL.CHEM
24362-6	Renal Function HCFA 2000 Panel	-	Pt	Ser/Plas	Panel		PANEL.CHEM
34566-0	Vital signs panel	-	Pt	^Patient	Panel		PANEL.VITALS
11488-4	Consultation note	Find	Pt	{Setting}	Doc	{Provider}	ATTACH.CLINRPT
34066-1	Boxed warning section	-	-	^FDA package insert	Doc		DOC.REF.
35511-5	Background information section	-	-	^Clinical trial protocol	Doc		DOC.REP.CTP
35660-0	Path report.final diagnosis section – text	Imp	Pt	Specimen	Doc		TUMRRGT
24534-0	Multisection	Find	Pt	Abdominal vessels	Doc	US.doppler	RAD

## 10 Standardized Assessment Measures

## ***10.1 Introduction***

The LOINC committee approved inclusion of standardized assessment measures (e.g., survey instruments) with version 1.0p. Representing the observations in these assessments within LOINC required a modest extension of the System axis to include aggregate units of analysis, such as “family”, and storage of additional attributes within the LOINC database. Bakken<sup>30</sup> provides a detailed description of the methodology for inclusion and evaluation into LOINC and the extensions to the LOINC axes.

The initial corpus of material represented in LOINC came from standardized nursing assessment instruments, including: Home Health Care Classification (HHC), Quality Audit Marker (QAM), Signs and Symptoms Checklist for Persons with HIV (HIV-SSC), Living with HIV, and the Omaha System. We have since expanded the content to cover standardized assessment instruments in many other domains.

## ***10.2 Consolidated Health Informatics endorsement***

As national interest in using standards for communicating the results of patient assessment instruments has increased, we have collaborated with members of the Consolidated Health Informatics (CHI) Disability Workgroup and many others to more fully develop the content and infrastructure to support patient assessment instruments. LOINC now contains full representations of CMS's Minimum Data Set version 2 (MDS) used in nursing homes, CMS's Outcome and Assessment Information Set (OASIS) used in home health care, and the Social Security Administration's Residual Functional Capacity (RFC) instrument.

Our work with CHI Disability Workgroup has led to the endorsement of Clinical LOINC as a CHI standard for federally-required assessment (i) questions and answers, and (ii) assessment form that include functioning and disability content. The recommendations of the CHI Disability Workgroup were endorsed by the NCVHS and subsequently the HHS Secretary.

## ***10.3 LOINC Representation***

The overall organization of the survey instruments are represented in LOINC using a nested panel structure consistent with the existing model for laboratory panels. LOINC codes are created for the individual questions/items within an instrument, as well as for the panels/groups of terms representing the hierarchical nature of the instrument. In addition, the database and RELMA program continue to be refined to support the definitional elements of the full instrument, including: computational and skip logic, help and contextual coding content, and structured answer lists.

### **10.3.1 Naming rules and conventions for survey instrument items in LOINC**

As we have expanded our work to represent more and varied patient assessment instruments within LOINC, we have evolved our model to capture more completely the relevant aspects of various forms. In particular, we have three fields that can capture the exact display of the question/item on the form in question. (For some instruments, it can be difficult to determine what exactly *is* the question text). The Component of a LOINC term represents the thing or attribute being measured, and is the default for capturing the item text. However, there are several reasons why the component may not be the exact item

text. The most important ones include: our LOINC naming conventions do not permit certain characters (e.g. "/" or "?") because of our internal "Part" parsing rules, that some aspects of the question are modeled in other parts of the formal LOINC name, or that there is some important aspect to the "thing being measured" (e.g. a lookback period of the last 7 days) that is not represented explicitly on the form for that particular item. In general, for purposes of displaying the item text as it appears on the instrument, one can follow this rule:

1. **SURVEY\_QUEST\_TEXT** in the LOINC table [if populated]. This field is populated when the variable/item is asked as a question. In some cases, the variable has both a question and a label. In these cases, the SURVEY\_QUEST\_TEXT field is populated with both, in the pattern of [Label].[Question text]. For example, for item J0300 on the MDS version 3 we have "Pain Presence. Ask resident: "Have you had pain or hurting at any time in the last 7 days?"
2. **DISPLAY\_NAME\_FOR\_FORM** in the FORM\_DATA table (RELMA) or FORMS table (CSV file export) [if populated]. This field provides an override display that is linked to the instance of the LOINC question code in a particular form. It allows for the same clinical concept to have slight presentation variances on different forms where those variances have no change in the concept meaning and accomodates instances where the LOINC naming conventions require some difference between the item and the LOINC Component. For example, an item might have the form label of "Body Mass Index (BMI)" but the LOINC Component would simply be "Body mass index".
3. **COMPONENT** in the LOINC table. This is the default display.

In addition to these fields, some LOINC codes used in survey instruments may have other LOINC name fields such as a Short Name, Long Common Name, or our newly created Consumer Name. These additional names may be useful in some contexts for these items, be we will still use the above rule to capture the item's representation in the instrument. Some of the original survey instruments modeled in LOINC may not follow this rule exactly, in part due to the fact that we did not have full survey representation model as we do presently. Ongoing work includes reviewing where modifications may be needed.

### 10.3.2 Structured answer lists

The questions/items in standardized assessment instruments often have highly specialized, fixed answer lists. In many contexts, it is the answer list that most completely defines the meaning of concept represented by the question. Additionally, because many of the answer choices are highly specialized, few are represented by existing codes in reference terminologies. For these reasons, we have created a structured representation of the answer lists for the questions in assessment instruments represented in LOINC. Individual answers are assigned a non-semantic identifier with a "LA" prefix and a mod-10 check digit. The answer codes LOINC assigns are unique by lexical string (ignoring capitalization), and by intention do not distinguish between strings that may have different meanings depending on their contextual use. The LA codes are placeholders for the structured answer lists, not intended as an external codes system or in the HL7 message.

## 11 Editorial Policies and Procedures

### *11.1 Concept orientation and LOINC name changes*

LOINC is a concept-based terminology, which means that it provides a way of naming classes of things that exist in the real world. Each concept (term) is given an identifier and a fully-specified name. Other attributes, including other names such as a Short Name and Long Common Name, are also provided in the LOINC database. The concept is anchored by the LOINC code, not by the particular strings in the formal name we happen to use to explain the code. It is certainly not possible to convey all of the subtleties that exist in the world with formal machinery alone, which is why we are also working very hard to include narrative text descriptions with each term that further elaborate and explain the concept.

In a complex, organic terminology like LOINC, name changes and modifications are unavoidable for many reasons. Since its inception, LOINC has maintained a set of editorial policies that guide our adherence to this concept-oriented ideal even as the terminology evolves over time. An over-arching policy is that we can change the name (i.e. the human-readable representation of the concept) in any way that does not change the meaning of the concept. In other words, a modification is allowable and valid if it is still an unambiguous reference to a class of things in the real world.

For example, two different numbering systems have been used to identify the serotypes of *Streptococcus pneumoniae*, which are important in gauging the coverage of polyvalent pneumonia vaccines. The U.S. system uses only numbers while the Danish system includes numbers and letters. For a period of time, LOINC term names were split and used a mixture of the Danish and U.S. identifiers, which was inconsistent and confusing. To clarify, we converted the few Danish serotype identifiers to their corresponding U.S. serotype identifiers (and included the Danish identifiers as synonyms). This was not a fundamental change to the underlying concept, but rather just the particular labels used to express it.

Not all situations are crystal clear, so our general policy is to seek as much input as is feasible, and typically such cases are brought for discussion to the LOINC Committee.

## ***11.2 Classification of LOINC term status***

LOINC development follows best practices for terminology system development by never reusing or deleting codes. If a LOINC term is identified as erroneous or a duplicate of a previous term it is flagged as “deprecated” in the database, but the record is not removed. Changes in concept status are made very judiciously. Prior to the LOINC version 2.31 release (June 2010), we identified such deprecated terms by populating the STATUS field of the database with “DEL” and wherever possible identified superseding concepts in the MAP\_TO field. Active (non-deprecated) records had no value (null) in the STATUS field. Based on new use cases and input from the LOINC community, LOINC 2.31 implements an expansion to that classification. The presently supported values for term status, with the definition and implications for use, are:

<b>ACTIVE</b>	Concept is active. Use at will.
<b>TRIAL</b>	Concept is experimental in nature. Use with caution as the concept and associated attributes may change.
<b>DISCOURAGED</b>	Concept is not recommended for current use. New mappings to this concept are discouraged; although existing may mappings may continue to be valid in context. Wherever possible, the superseding concept is indicated in the MAP_TO field of the database and should be used instead.
<b>DEPRECATED</b>	Concept is deprecated. Concept should not be used, but it is retained in LOINC for historical purposes. Wherever possible, the superseding concept is indicated in the MAP_TO field of the database and should be used both for new mappings and updating existing implementations.

Furthermore, we have added two new fields:

<b>STATUS_REASON</b>	Classification of the reason for concept status. This field will be Null for ACTIVE concepts, and optionally populated for terms in other statuses where the reason is clear. DEPRECATED or DISCOURAGED terms may take values of: AMBIGUOUS, DUPLICATE, or ERRONEOUS.
<b>STATUS_TEXT</b>	Explanation of concept status in narrative text. This field will be Null for ACTIVE concepts, and optionally populated for terms in other statuses.

Our initial implementation of these new concept status values populated the STATUS field with “ACTIVE” or “DEPRECATED” based on their existing status and have identified a limited set of terms that have been designated “DISCOURAGED” or “TRIAL”.

The principal reason identifying terms as DISCOURAGED is where we have strong inclinations that a particular term is no longer valid given current practice. For example, we have flagged as DISCOURAGED several lutropin terms with Properties consistent with mass or molar units because all lutropin sources that we could reach report concentrations in international units (IU) per volume and the drug lutropin is prescribed in terms of international units per volume. To avoid help confusion on the part of mappers, the DISCOURAGED status steers them away from these terms to the more likely candidates.

The principal reason for identifying terms as TRIAL is a very constrained circumstance, such as when the source of the term is still equivocating. This has been illustrated in our work to create a LOINC representation of federally-required patient assessment instruments. Here, the item meaning is defined in the context of use within that instrument. We have been fortunate to work with assessment developers in the early stages of instrument development. This is advantageous because it enables the codes to be included in data specifications and documents as they are developed, but we are in the position of creating codes and names for data elements whose attributes are still in flux. As the instrument evolves, the specific representation of the item (question) or answer options on the form may change. Ultimately, the representation will be settled by an authoritative body (such as CMS) and they are intended for use in one context – the official release of the instrument. Identifying these terms as TRIAL allows us to include them in the public distribution while clearly flagging their “pending” status. Once the final concept representation has been determined, terms initially labeled TRIAL would be reclassified as ACTIVE (or perhaps in rare circumstances DEPRECATED or even rarer DISCOURAGED).

### ***11.3 Concept persistence and term deprecation***

LOINC codes are never reused or deleted, and the concept meaning is persistent over time despite the fact that there may be modifications to the name (as described above). If we discover that a LOINC term's meaning is a duplicate of another existing term or it is somehow erroneous, it will be given a status of DEPRECATED but not removed from the database.

In the past, when we encountered duplicate terms (i.e. terms that had different names but meant the same thing), our general policy was to deprecate the newest term. Many times, this convention worked well because the older term was more likely to have been incorporated into user's systems through mappings, etc. However, this was not always the case. Sometimes, the new term had the clearer, more recognizable label and thus it was most likely the most mapped-to term.

Therefore, our current policy is to make the change to require (in our estimation) the least amount of re-mapping for existing users.

## Appendix A - LOINC Database Structure

Table 28: LOINC Database Structure			
Field Name	Type	Width	Description
1. LOINC_NUM	Text	7	The unique LOINC Code is a string in the format of nnnnn-n.
2. COMPONENT	Text	255	First major axis-component or analyte
3. PROPERTY	Text	30	Second major axis-property observed (e.g., mass vs. substance)
4. TIME_ASPCT	Text	15	Third major axis-timing of the measurement (e.g., point in time vs 24 hours)
5. SYSTEM	Text	100	Fourth major axis-type of specimen or system (e.g., serum vs urine)
6. SCALE_TYP	Text	30	Fifth major axis-scale of measurement (e.g., qualitative vs. quantitative)
7. METHOD_TYP	Text	50	Sixth major axis-method of measurement
8. RELAT_NMS	Text	254	This field is no longer being maintained. It has been replaced by RelatedNames2
9. CLASS	Text	20	An arbitrary classification of the terms for grouping related observations together. The current classifications are listed in Table 29. We present the database sorted by the class field within class type (see field 23). Users of the database should feel free to re-sort the database in any way they find useful, and/or to add their own classifying fields to the database.  The content of the laboratory test subclasses should be obvious from the subclass name.
10. SOURCE	Text	8	This is for our internal use and should be ignored by database users.
11. DT_LAST_CH	Text	8	Date last changed, in the format YYYYMMDD
12. CHNG_TYPE	Text	3	Change Type Code DEL = delete (deprecate) ADD = add NAM = change to Analyte/Component (field #2); MAJ = change to name field other than #2 (#3 - #7); MIN = change to field other than name UND = undelete
13. COMMENTS	Memo	-	Free-text comments relating to the test result.
14. ANSWERLIST	Memo	-	The contents of this field have been moved to new structured lists which can be viewed in the details screen using RELMA.
15. STATUS	Text	11	ACTIVE = Concept is active. Use at will. TRIAL = Concept is experimental in nature. Use with caution as the concept and associated attributes may change. DISCOURAGED = Concept is not recommended for current use. New mappings to this concept are discouraged; although existing mappings may continue to be valid in context. Wherever possible, the superseding concept is indicated in the MAP_TO field of the database and should be used instead. DEPRECATED = Concept is deprecated. Concept should not be used, but it is retained in LOINC for historical purposes. Wherever possible, the superseding concept is indicated in the MAP_TO field of the database and should be used both for new mappings and updating existing implementations.
16. MAP_TO	Text	7	Used when a field has been dropped from the active database (by entering "DEL" in the Status field) because it has been replaced by an updated term. In those cases, MAP_TO contains the LOINC code of the new term that should be used.
17. SCOPE	Text	20	Not currently used.
18. CONSUMER_NAME	Text	255	An experimental (beta) consumer friendly name for this item. The intent is to provide a test name that health care consumers will recognize; it will be similar to the names that might appear on a lab report and is not guaranteed to be unique

			because some elements of the LOINC name are likely to be omitted. We will continue to modify these names in future release, so do not expect it to be stable (or perfect). Feedback is welcome.
19. IPCC_UNITS	Text	30	Units have been moved to a new structure that can be viewed in the details screen in RELMA.
20. REFERENCE	Memo	-	Contains references to medical literature, product announcements, or other written sources of information on the test or measurement described by the LOINC record.
21. EXACT_CMP_SY	Text	50	Exact core component synonym: This field contains an exact synonym for the "core component" of the LOINC component name. We have included the mixed case and "superscript" form of blood bank and HLA antigens (e.g., Lua) here. As there is no ASCII representation for superscript letters, we use the hat (^) to signify superscripts in this field. (e.g., if the core component is represented as "L little u super little a" in the LOINC component/analyte name field, it is represented in the Exact Core Synonym field as Lu^a.)
22. MOLAR_MASS	Text	13	Molecular weights: This field contains the molecular weights of chemical moieties when they are provided to us. This release contains values kindly contributed by IUPAC.
23. CLASSTYPE	Int	2	1=Laboratory class; 2=Clinical class; 3=Claims attachments; 4=Surveys
24. FORMULA	Text	255	Regression equation details for many OB.US calculated terms.
25. SPECIES	Text	20	Codes detailing which non-human species the term applies to. If blank, "human" is assumed.
26. EXMPL_ANSWERS	Memo	-	For some tests and measurements, we have supplied examples of valid answers, such as "1:64", "negative @ 1:16", or "55".
27. ACSSYM	Memo	-	Chemical name synonyms, alternative name synonyms, and chemical formulae supplied by the Chemical Abstract Society.
28. BASE_NAME	Text	50	Chemical base name from CAS
29. FINAL	Text	1	Internal LOINC use field
30. NAACCR_ID	Text	20	Maps to North American Association of Central Cancer Registries Identification Number
31. CODE_TABLE	Text	10	Examples on CR0050 Cancer Registry
32. SETROOT	Yes/No	1	Currently used for claims attachments. Yes in this field signifies that this record is the root of a set of LOINC codes. 0=No, 1=Yes
33. PANELEMENTS	Memo	-	This field is no longer being maintained. See "Viewing LOINC Details" in RELMA.
34. SURVEY_QUEST_TXT	Memo	-	Verbatim question from the survey instrument
35. SURVEY_QUEST_SRC	Text	50	Exact name of the survey instrument and the item/question number
36. UNITSREQUIRED	Text	1	Y/N field that indicates that units are required when this LOINC is included as an OBX segment in a HIPAA attachment
37. SUBMITTED_UNITS	Text	30	Units as received from person who requested this LOINC term.
38. RELATEDNAMES2	Memo	-	This is a new field introduced in version 2.05. It contains synonyms for each of the parts of the fully specified LOINC name (component, property, time, system, scale, method). It replaces #8, Relat NMS.
39. SHORTNAME	Text	40	Introduced in version 2.07, this field is a concatenation of the fully specified LOINC name. The field width may change in a future release.
40. ORDER_OBS	Text	15	Defines term as order only, observation only, or both. A fourth category, Subset, is used for terms that are subsets of a panel but do not represent a package that is known to be orderable we have defined them only to make it easier to maintain panels or other sets within the LOINC construct.
41. CDISC_COMMON_TESTS	Text	1	"Y" in this field means that the term is a part of subset of terms used by CDISC in clinical trials.
42. HL7_FIELD_SUBFIELD_ID	Text	50	A value in this field means that the content should be delivered in the named field/subfield of the HL7 message. When NULL, the data for this data element should be sent in an OBX segment with this LOINC code stored in OBX-3 and

			with the value in the OBX-5.
43. EXTERNAL_COPYRIGHT_NOTICE	Memo	-	External copyright holders copyright notice for this LOINC code.
44. EXAMPLE_UNITS	Text	255	This field is populated with a combination of submitters units and units that people have sent us. Its purpose is to show users representative, but not necessarily recommended, units in which data could be sent for this term.
45. INPC_PERCENTAGE	Number	-	Percent of overall 13 month result volume within the INPC (circa 2006). A record with a value > 0 indicates that this code is one of the common tests that comprise > 99% of the total INPC result volume. See also PMID: 18693941 .
46. LONG_COMMON_NAME	Text	255	This field contains the LOINC term in a more readable format than the fully specified name. The long common names have been created via a table driven algorithmic process. Most abbreviations and acronyms that are used in the LOINC database have been fully spelled out in English.
47. HL7_V2_DATATYPE	Text	255	HL7 version 2.x data type that would be sent in OBX-2 when this data is delivered in an HL7 message.
48. HL7_V3_DATATYPE	Text	255	HL7 version 3.0 data type that is compatible with this LOINC code.
49. CURATED_RANGE_AND_UNITS	Memo	-	A curated list of normal ranges and associated units (expressed as near UCUM codes) for physical quantities and survey scores. Intended as tailorable starter sets for applications that use LOINC forms as a way to capture data. Units are separated from normal ranges by XXX and sets of normal range/units pairs are separated by YYY.
50. DOCUMENT_SECTION	Text	255	Classification of whether this LOINC code can be used a full document, a section of a document, or both. This field was created in the context of HL7 CDA messaging, and populated in collaboration with the HL7 Structured Documents Technical Committee.
51. DEFINITION_DESCRIPTION_HELP	Memo	-	Field that is similar to the current COMMENTS field that tells users what the LOINC term is and what to do with it.
52. EXAMPLE_UCUM_UNITS	Text	255	The Unified Code for Units of Measure (UCUM) is a code system intended to include <i>all</i> units of measures being contemporarily used in international science, engineering, and business. ( <a href="http://www.unitsofmeasure.org">www.unitsofmeasure.org</a> ) This field contains example units of measures for this term expressed as UCUM units.
53. EXAMPLE_SI_UCUM_UNITS	Text	255	The Unified Code for Units of Measure (UCUM) is a code system intended to include <i>all</i> units of measures being contemporarily used in international science, engineering, and business. ( <a href="http://www.unitsofmeasure.org">www.unitsofmeasure.org</a> ) This field contains example units of measures for this term expressed as SI UCUM units.
54. STATUS_REASON	Text	9	Classification of the reason for concept status. This field will be Null for ACTIVE concepts, and optionally populated for terms in other status where the reason is clear. DEPRECATED or DISCOURAGED terms may take values of: AMBIGUOUS, DUPLICATE, or ERRONEOUS.
55. STATUS_TEXT	Memo	-	Explanation of concept status in narrative text. This field will be Null for ACTIVE concepts, and optionally populated for terms in other status.
56. SPECIAL_EXPLANATION	Memo	-	Detailed explanation about special changes to the term over time.

## Appendix B - Classes

Table 29: Classes

Table 29a: Clinical Term Classes	
Abbreviation	Clinical Term Class
ART	Antiretroviral therapy
BDYCRC.ATOM	Body circumference atomic
BDYCRC.MOLEC	Body circumference molecular
BDYHGT.ATOM	Body height atomic
BDYHGT.MOLEC	Body height molecular
BDYSURF.ATOM	Body surface atomic
BDYTMP.ATOM	Body temperature atomic
BDYTMP.MOLEC	Body temperature molecular
BDYTMP.TIMED.MOLEC	Body temperature timed molecular
BDYWGT.ATOM	Body weight atomic
BDYWGT.MOLEC	Body weight molecular
BP.ATOM	Blood pressure atomic
BP.CENT.MOLEC	Blood pressure central molecular
BP.MOLEC	Blood pressure molecular
BP.PSTN.MOLEC	Blood pressure positional molecular
BP.TIMED.MOLEC	Blood pressure timed molecular
BP.VENOUS.MOLEC	Blood pressure venous molecular
CARD.US	Cardiac ultrasound (was US.ECHO)
CLIN	Clinical NEC (not elsewhere classified)
DENTAL	Dental
DEVICES	Medical devices
DOC.ADMIN	Administrative documents
DOC.CLINRPT	Clinical report documentation
DOC.EPSOS	Smart Open Services for European Patients (epSOS) documents
DOC.MISC	Miscellaneous documentation
DOC.PUBLICHEALTH	Public health documentation
DOC.QUALITY	Quality documents
DOC.REF	Referral documentation
DOC.REF.CTP	Clinical trial protocol document
DOCUMENT.REGULATORY	Regulatory documentation
ED	Emergency (DEEDS)
EKG.ATOM	Electrocardiogram atomic
EKG.IMP	Electrocardiogram impression
EKG.MEAS	Electrocardiogram measures
ENDO.GI	Gastrointestinal endoscopy
EYE	Eye
EYE.CONTACT_LENS	Ophthalmology contact lens
EYE.GLASSES	Ophthalmology glasses: Lens manufacturer (LM) & Prescription
EYE.HETEROPHORIA	Ophthalmology heterophoria
EYE.OCT	Ophthalmology Optical Coherence Tomography (OTC)
EYE.PX	Ophthalmology physical findings
EYE.REFRACTION	Ophthalmology refraction
EYE.RETINAL_RX	Ophthalmology treatments
EYE.TONOMETRY	Ophthalmology tonometry

EYE.US	Ophthalmology ultrasound
EYE.VISUAL_FIELD	Ophthalmology visual field
FUNCTION	Functional status (e.g., Glasgow)
H&P.HX	History
H&P.PX	Physical
H&P.SURG PROC	Surgical procedure
HEMODYN.ATOM	Hemodynamics anatomic
HEMODYN.MOLEC	Hemodynamics molecular
HRTRATE.ATOM	Heart rate atomic
HRTRATE.MOLEC	Heart rate molecular
HRTRATE.TIMED.MOL	Heart rate timed molecular
IO.TUBE	Input/Output of tube
IO_IN.ATOM	Input/Output atomic
IO_IN.MOLEC	Input/Output molecular
IO_IN.SUMMARY	Input/Output summary
IO_IN.TIMED.MOLEC	Input/Output timed molecular
IO_IN_SALTS+CALC	Input/Output electrolytes and calories
IO_OUT.ATOM	Input/Output atomic
IO_OUT.MOLEC	Input/Output molecular
IO_OUT.TIMED.MOLE	Input/Output timed molecular
NEONAT	Neonatal measures
OB.US	Obstetric ultrasound
OBGYN	Obstetric/Gynecology
PANEL.ART	Antiretroviral therapy order set
PANEL.BDYTMP	Body temperature order set
PANEL.BP	Blood pressure order set
PANEL.CARDIAC	Cardiac studies order set
PANEL.CV	Cardiovascular order set
PANEL.DEVICES	Medical devices order set
PANEL.DOC	Documents panels
PANEL.DOC.CLINRPT	Clinical report documentation set
PANEL.ED	Emergency (DEEDS) order set
PANEL.EYE	Ophthalmology panels
PANEL.FUNCTION	Function order set
PANEL.H&P	History & Physical order set
PANEL.IO	Input/Output order set
PANEL.NEONAT	Neonatal measures order set
PANEL.OB.US	Obstetrical order set
PANEL.PATIENT SAFETY	Patient safety order set
PANEL.PHENX	PhenX Panel
PANEL.PHR	Public health record order set
PANEL.RAD	Radiology order set
PANEL.TUMRRGT	Tumor registry order set
PANEL.US.URO	Urology ultrasound order set
PANEL.VACCIN	Vaccination order set
PANEL.VITALS	Vital signs order set
PATH.PROTOCOLS.BRST	Pathology protocols - breast
PATH.PROTOCOLS.GENER	Pathology protocols - general
PATH.PROTOCOLS.PROST	Pathology protocols - prostate

PATH.PROTOCOLS.SKIN	Pathology protocols - skin
PATIENT SAFETY	Patient safety
PHENX	PhenX
PUBLICHEALTH	Public Health
PULM	Pulmonary ventilator management
RAD	Radiology
RESP.ATOM	Respiratory atomic
RESP.MOLEC	Respiratory molecular
RESP.TIMED.MOLEC	Respiratory timed molecular
SKNFLD.MOLEC	Skinfold measurements molecular
TRNSPLNT.ORGAN	Organ transplant
TUMRRGT	Tumor registry (NAACCR)
US.URO	Urological ultrasound
VACCIN	Vaccinations
VOLUME.MOLEC	Volume (specimen) molecular

<b>Table 29b: Laboratory Term Classes</b>	
<b>Abbreviation</b>	<b>Laboratory Term Class</b>
ABXBACT	Antibiotic susceptibilities
ALLERGY	Response to antigens
BLDBK	Blood bank
CELLMARK	Cell surface models
CHAL	Challenge tests
CHALSKIN	Skin challenge tests
CHEM	Chemistry
COAG	Coagulation study
CYTO	Cytology
DRUG/TOX	Drug levels & Toxicology
DRUGDOSE	Drug dose (for transmitting doses for pharmacokinetics)
FERT	Fertility
HEM/BC	Hematology (coagulation) differential count
HL7.GENETICS	Clinical genetic report
HL7.CYTOGENETICS	Clinical cytogenetic report
HLA	HLA tissue typing antigens and antibodies
HPA	HPA typing
LABORDERS	Laboratory order codes
MICRO	Microbiology
MISC	Miscellaneous
MOLPATH	Molecular pathology
MOLPATH.DEL	Gene deletion
MOLPATH.MISC	Gene miscellaneous
MOLPATH.MUT	Gene mutation
MOLPATH.REARRANGE	Gene rearrangement
MOLPATH.TRINUC	Gene trinucleotide repeats
MOLPATH.TRISOMY	Gene chromosome trisomy
MOLPATH.TRNLOC	Gene translocation

NR STATS	Normal range statistics
PANEL.ABXBACT	Susceptibility order sets
PANEL.ALLERGY	Allergy order set
PANEL.BLDBK	Blood bank order set
PANEL.CELLMARK	Cell marker order sets
PANEL.CHAL	Challenge order set
PANEL.CHEM	Chemistry order set
PANEL.COAG	Coagulation order set
PANEL.DRUG/TOX	Drug level & Toxicology order set
PANEL.FERT	Fertility testing order set
PANEL.HEDIS	Healthcare Effectiveness Data and Information Set order set
PANEL.HEM/BC	Hematology & blood count order set
PANEL.HL7.GENETICS	HL7 genetics panel
PANEL.HL7.CYTOGENETICS	HL7 cytogenetics panel
PANEL.HLA	HLA order set
PANEL.HPA	HPA order set
PANEL.MICRO	Microbiology order set
PANEL.MISC	Miscellaneous order set
PANEL.MOLPATH	Molecular pathology order set
PANEL.OBS	Obstetrics order set
PANEL.SERO	Serology order set
PANEL.UA	Urinalysis order set
PATH	Pathology
SERO	Serology (antibodies and most antigens except blood bank and infectious agents)
SPEC	Specimen characteristics
UA	Urinalysis

**Table 29c: Attachment Term Classes**

<b>Abbreviation</b>	<b>Attachment Term Class</b>
ATTACH	Attachment
ATTACH.AMB	Ambulance attachment
ATTACH.CARD	Cardiac attachment
ATTACH.CLINRPT	Clinical report attachment
ATTACH.CPHS	Children's Preventative Health System attachment
ATTACH.ED	Emergency department attachment
ATTACH.GENERAL	General attachment
ATTACH.GI	Gastrointestinal attachment
ATTACH.LAB	Laboratory attachment
ATTACH.MEDS	Medication attachment
ATTACH.MODIFIER	Modifier attachment
ATTACH.OBS	Obstetrics attachment
ATTACH.REHAB	Rehabilitation attachment
ATTACH.REHAB.ABUSE	Alcohol/Substance abuse rehabilitation attachment
ATTACH.REHAB.CARDIAC	Cardiac rehabilitation attachment

ATTACH.REHAB.NURS	Specialized nursing attachment
ATTACH.REHAB.OT	Occupational therapy attachment
ATTACH.REHAB.PSYCH	Psychiatric rehabilitation attachment
ATTACH.REHAB.PT	Physical rehabilitation attachment
ATTACH.REHAB.PULM	Pulmonary rehabilitation attachment
ATTACH.REHAB.RT	Respiratory rehabilitation attachment
ATTACH.REHAB.SOCIAL	Medical social work attachment
ATTACH.REHAB.SPEECH	Speech therapy rehabilitation attachment
ATTACH.RESP	Respiratory attachment

<b>Table 29d: Survey Term Classes</b>	
<b>Abbreviation</b>	<b>Survey Term Class</b>
PANEL.SURVEY.BIMS	Brief Interview for Mental Health Status (BIMS) set
PANEL.SURVEY.CARE	Continuity Assessment Record and Evaluation (CARE) set
PANEL.SURVEY.CAM	Confusion Assessment Method (CAM) set
PANEL.SURVEY.GDS	Geriatric Depression Scale (GDS) set
PANEL.SURVEY.HHCC	Home Health Care Classification set
PANEL.SURVEY.HIV-SSC	Signs and Symptoms checklist for persons living with HIV set
PANEL.SURVEY.HOWRU	howRU outcomes instrument set
PANEL.SURVEY.LIV-HIV	Living with HIV set
PANEL.SURVEY.MDS	Minimum Data Set for Nursing Home Resident Assessment and Care Screening set
PANEL.SURVEY.MFS	Morse Fall Scale set
PANEL.SURVEY.NMMDS	Nursing Management Minimum Data set
PANEL.SURVEY.OASIS	Outcome and Assessment Information Survey set
PANEL.SURVEY.OMAHA	OMAHA survey set
PANEL.SURVEY.PHQ9	Patient Health Questionnaire PHQ-9 set
PANEL.SURVEY.PROMIS	Patient Reported Outcomes Measurement System set
PANEL.SURVEY.QAM	Quality Audit Marker set
PANEL.SURVEY.QRDA	Quality Health Reporting Document Architecture set
PANEL.SURVEY.RFC	Residual Functional Capacity set
PANEL.TIMP	Test of Infant Motor Performance set
PANEL.SURVEY.USSGFHT	United States Surgeon General Family Health Tool set
SURVEY.CARE	Continuity Assessment Record and Evaluation (CARE) survey
SURVEY.GDS	Geriatric Depression Scale (GDS) survey
SURVEY.HOWRU	howRU outcomes instrument survey
SURVEY.MDS	Minimum Data Set for Nursing Home Resident Assessment and Care Screening survey
SURVEY.MFS	Morse Fall Scale survey
SURVEY.NMMDS	Nursing Management Minimum Data survey
SURVEY.NURSE.HHCC	Home Health Care Classification survey
SURVEY.NURSE.HIV-SSC	Signs and Symptoms checklist for persons living with HIV survey
SURVEY.NURSE.LIV-HIV	Living with HIV survey
SURVEY.NURSE.OMAHA	OMAHA survey
SURVEY.NURSE.QAM	Quality Audit Marker survey
SURVEY.OASIS	Outcome and Assessment Information Survey
SURVEY.PHQ	Patient Health Questionnaire
SURVEY.QRDA	Quality Health Reporting Document Architecture survey

SURVEY.PROMIS	Patient Reported Outcomes Measurement System survey
SURVEY.RFC	Residual Functional Capacity survey
SURVEY.USSGFHT	United States Surgeon General Family Health Tool survey
TIMP	Test of Infant Motor Performance survey

## Appendix C - Calculating Mod 10 Check Digits

The algorithm for calculating a Mod 10 check digit is as follows:

### Instructions

	<b>Example</b>
1. Using the number 12345, assign positions to the digits, from right to left.	1st = 5 2nd = 4 3rd = 3 4th = 2 5th = 1
2. Take the odd digit positions counting from the right (1st, 3rd, 5th, etc.)	531
3. Multiply by 2.	1062
4. Take the even digit positions starting from the right (2nd, 4th, etc.).	42
5. Append (4) to the front of the results of (3).	421062
6. Add the digits of (5) together.	$4+2+1+0+6+2 = 15$
7. Find the next highest multiple of 10.	20
8. Subtract (6) from (7).	
Thus, 5 is the Mod 10 check digit for 12345.	$20 - 15 = 5.$

## **Appendix D - Procedure for Submitting Additions/Changes to the Database**

### **Introduction**

The Regenstrief Institute receives two kinds of requests for additions:

- (1) The first kind of request deals with (a) an entirely new kind of measurement, e.g., DNA sequencing or (b) the use of LOINC codes in manners that have not been agreed upon by the LOINC committee, e.g., the definition of terms to accommodate the organism 1, organism 2, etc., structures that are present in many laboratory databases.
- (2) Other requests are variations on observations that are already in the database. For example, we have a term for a particular test result with serum as the specimen (system) and a user requests an identical term for a specimen of gastric contents. Provided that the requestor followed the rules given below and the number of terms requested at a given time is modest, we will try to respond to these kinds of requests quickly.

The Institute will only be able to respond quickly to such requests if the requestor provides us with clear information about the new terms, as detailed below in Table 30, which defines the content that we need to determine whether a submitted code requires a new LOINC code assignment or not. Before sending a request, make sure that you have, at a minimum, provided information about the component, property, timing aspect, system, scale, method and units of measure (when appropriate). It is also very useful for us to receive example results or answers lists of the test/observation that is being requested. This information enables us to verify the property, scale, and method.

You have the option of either submitting a file produced solely by you or one generated on your behalf via the RELMA program. Regardless of which option you choose, your submission file must be sent to the Regenstrief Institute in one of three file formats.

- The preferred format (and the one that RELMA will produce on your behalf) is a Microsoft Access database (mdb).
- The second format is a Microsoft Excel spreadsheet (xls).
- The final format is a tab delimited ASCII file (txt).

The example file and field descriptions below will aid you in creating a submission file from scratch (without the aid of the RELMA program).

### **A Few Notes before Proceeding**

The terms “addition”, “requested term” and “proposed LOINC” are synonymous. All of these terms refer to a concept created by a user that will be or has been submitted to the Regenstrief Institute for consideration as an addition to the LOINC database.

Please note that we tend to avoid the use of methods for chemistry tests. We will not routinely accept requests for method-specific chemistry tests. Only in very special circumstances will we distinguish among analytic methods in chemistry. We do distinguish microbiology, serology, and coagulation tests by method type. Even here, however, we do not distinguish every variation in methods. Look in the body of this guide for information about the kinds of distinctions that we make.

If you find a test in the database that you believe is wrong, please send us a letter or email ([loinc@regenstrief.org](mailto:loinc@regenstrief.org)) calling attention to the term and the reason you think it is wrong, (e.g., not using

the standard nomenclature, typographical error, system of serum when it is only valid when performed on plasma, duplicate of some other concept in the database, etc. We welcome all input from users.

Note that our policy is to allow both method-vague (no method) as well as method-specific measures in serology (measures of Ab and Ag), and in antibiotic susceptibility testing.

Please pay special attention to requests for submissions that include the system of serum or plasma alone. For most chemical analyses there is no important clinical difference between the values obtained from serum and those obtained from plasma, and we would like to represent them in the database as Ser/Plas to indicate our indifference to the distinction. Unfortunately, many requestors of new terms define their request in terms of the one that they happen to use (e.g., serum or plasma) without telling us that the measure can really be done on either serum or plasma. Most such requests should be for Ser/Plas as the system (sample). If the measurement **MUST** be done on either serum or plasma, please scientifically justify your request; otherwise you will greatly delay our response to your submission.

When possible, please provide a full description of the test, its purpose, and procedure. (A copy of vendor's test kit descriptive material or a copy from a textbook describing the procedure and its purpose would be very helpful.) We often will require a committee discussion to decide how to represent new subject matter, so response times will be slower.

The requestors also need to supply some evidence that they are familiar with the database and that they are sure the term is not already represented in LOINC. The major work these requests generate is the effort to be sure the observation is not already in the database. We can perform this service if the requestors have done most of this work themselves. For this reason, we request that you identify the LOINC term that is closest to your request and to flag the difference between the requested test and the existing test. That is, when a new observation is only a variation on an old one, use an existing LOINC observation as the template, change the part that is different in the new term and indicate that difference.

### An Example Submission and Definition of the Submission File

An example submission (which, because of space limitations, includes columns for only the first few fields) appears below. Real submissions should have columns for all of the items listed in Table 29. Additional details are provided in the sections on creating Access Database and Excel submissions presented later in this appendix.

Row#	Your test ID	Analyte/Component	Property	Time	System	Scale	Method	Related	Etc
1	G23	Glucose^90M post 50g lactose	MCnc	Pt	Urine	Ord	Test strip		
2	C47	Coproporphyrin 1 isomer	MRat	24H	Urine	Qn			
3	I98	Indican	MRat	24H	Urine	Qn			
4	T51	Thyroxine.free	MCnc	Pt	Urine	Qn			

The following table contains a description of the fields contained in the submit.mdb template. These fields should be present in the submission file you submit to Regenstrief. Only some of the fields need to be populated with data as noted below.

Field Name	Type	Width	Description
1. S_ROW	Long	4	Row number of this term in submitter's file.
2. S_LOCAL_CD	Text	50	The submitter's local code used to identify the test/observation in the submitter's master file.
3. S_COMPO	Text	150	Submitter's Analytes/Component. Mandatory. (User Guide 2.2)
4. S_PROP	Text	30	Submitter's Kind of Property. Mandatory – but we can help if you provide enough details. (User Guide 2.3)
5. S_TIME	Text	15	Submitter's Time Aspect. Mandatory. (User Guide 2.4)
6. S_SYS	Text	100	Submitter's System/Sample Type. Mandatory. (User Guide 2.5)
7. S_SCALE	Text	30	Submitter's Type of Scale. Mandatory. (User Guide 2.6)
8. S_METH	Text	50	Submitter's Type of Method. If required. (User Guide 2.7)
9. S_REL_NAM	Text	254	Submitter's Related Names. Strongly recommended. Common names, acronyms or synonyms.
10. S_LOINC	Text	10	Submitter's LOINC number. Strongly recommended. This is the LOINC number that is similar, but not the same as, the submitter's test.
11. S_RESULTS	Memo	-	Submitter's Example Results. Recommended for new analytes. As reported by your lab.
12. S_UNITS	Text	30	Submitter's Example Units. Mandatory for quantitative results. As reported by your lab.
13. S_SPECIES	Text	20	To be used for veterinary term submissions.
14. S_ID	Text	50	If the submitter includes a reference code ID for each unique submission to LOINC, record that ID here, and this will be returned with questions or an assigned LOINC number on a returned file.
15. S_COMMENT	Memo	-	Comments the submitter may wish to pass to RI when needed.

### Creating a Submission Using Microsoft Access

A blank Microsoft Access database template (.mdb) is available at <http://loinc.org/submissions>.




Figure 1. Submission Created with Microsoft Access

### Creating a Submission Using Microsoft Excel

If you choose to create your submission using Microsoft Excel, you should use the field names as specified in Table 29. Failure to do so may result in delays in processing the submission. Figure 2 below

shows an example of what an Excel submission template looks like.  
An Excel submission template (.xls) can be downloaded at <http://loinc.org/submissions>.

	A	B	C	D	E	F	G	H	I
1	S_ROW	S_LOCAL_CD	S_COMPO	S_PROP	S_TIME	S_SYS	S_SCALE	S_METH	S_REL_NAM
2									
3	1	G23	GLUCOSE*9 0M POST 50G LACTOSE PO	MCNC	T	UR	ORD	TEST STRIP	CHAL
4									
5	2	C47	COPROPORP HYRIN 1 ISOMER	MRAT	24H	UR	QN		CHEM
6									
7	3	I98	INDICAN	MRAT	24H	UR	QN		CHEM
8									
9	4	T51	THYROXINE FREE	MCNC	PT	UR	ORD		
10									
11									
12									
13									
14									

Figure 2. An Example Excel Submission (first 9 fields only)

**CAUTION:** Please take note of the field size indicated in Table 29. Upon receipt of your submission, we will copy the submission data into a Microsoft Access database as defined above in Table 29. If the cells in your Excel submission contain too many characters, some data may be lost in the conversion process.

### Creating a Submission Using a Tab-Delimited ASCII Text File

If you choose to send your submission in a tab-delimited ASCII text file format, please use the following format:

```
S_ROW|S_LOCAL_CD|S_COMPO|S_PROP|S_TIME|S_SYS|S_SCALE|S_METH|S_REL_NAM|S_LO
INC|S_RESULTS|S_UNITS|S_ID|S_SPECIES|S_COMMENT<CRLF>
```

Each field is separated from the other by a Tab character. That is, each vertical bar above would actually be a Tab character (i.e., an ASCII 9). A carriage-return/line-feed pair (i.e., the <CRLF> above) terminates each line. Therefore, each <CRLF>-terminated line in the ASCII file becomes a submission record. Note that the field lengths presented in 29 still apply to ASCII file submissions because upon receipt of your submission we copy the ASCII file data you submit into an Access database of the form described above.

**Using the previous example, one line might appear as:**

1|G23|GLUCOSE^90m POST 50g LACTOSE PO|MCNC|PT|UR|ORD|TEST STRIP|||6762-9||MG/DL||

where the vertical bars represent the Tab character. Notice that two vertical bars appear between “TEST STRIP” and “6762-9”. In this example, this means that the related names field is empty (i.e., a null field value). The example also shows that fields S\_RESULTS, S\_ID, and S\_COMMENT are also empty. Without the empty field, the field information would get out of sync and it would appear that the related names for this submission were actually the closest LOINC number for the submission (i.e., “6762-9”). Therefore, the ordering of the fields and the use of the Tab character to delimit the fields is very important.

In Figure 3 below, an example submission file is shown with the actual tab characters in lieu of the vertical bars used above as illustrations. Please note that the first row contains the field names described in Table 29. Also note that the tab characters are invisible to the human eye and make the text appear chaotic (this is one reason we recommend the use of Microsoft Access for the creation of submission files).

S_ROW	S_LOCAL_CD	S_COMPO	S_PROP	S_TIME	S_SYS	S_SCALE	S_METH	S_REL_NAM
1	G23	GLUCOSE^90m POST 50g LACTOSE PO	MCNC	T	UR	ORD	TEST STRIP	
2	C47	COPROPORPHYRIN 1 ISOMER MRAT	24H	UR	QN			
3	I98	INDICAN MRAT	24H	UR	QN			
4	T51	THYROXINE.FREE	MCNC PT	UR	ORD			

Figure 3. Example Submission Using ASCII Tab-Delimited File

## Generating a Submission Using RELMA

The RELMA program can aid you in creating submissions by allowing you to create, manage and store submission terms in a way that is similar to how the program creates, manages and stores local working sets. With RELMA, you can create terms for submission over time and submit groups of terms in batches. The program will track when the term was created and the date when you submitted the term. The program will help you organize the terms that you create and it will automate the process of creating the submission files.

For detailed instructions on how to use the RELMA propose a term feature, please see Appendix B of the RELMA manual which can be downloaded at <http://loinc.org/downloads/files/RELMAManual.pdf>.

## Sending Submission Files to Regenstrief Institute

Regardless of the file type, please email to [submissions@loinc.org](mailto:submissions@loinc.org). Note that you may have to zip the file. If you do not have access to email, you may copy this file onto a CD and mail it to:

Kathy Mercer  
 The Regenstrief Institute, Inc.  
 Health Information and Translational Sciences Bldg. (HITS)  
 410 West 10th Street, Suite 2000

Indianapolis, IN 46202

Within a day or two of receipt of your file, you will receive a confirmation email and the submission process will be underway. You may receive additional communication from Regenstrief with requests for further information if required. Once the submission process has completed, you will receive files containing your requested codes.

## Appendix E - Examples for LOINC Property Matching

1. **Content (CNT)** Like concentration except that volume in the denominator is replaced by mass. By extension:

CCnt Catalytic Content, catalytic activity of a component per unit mass of a sample (system).  
24048-1|Alpha galactosidase:CCnt:Pt:Fib:Qn

MCnt Mass Content, mass of component per unit mass of a sample (system).  
9435-9|Isopropanol:MCnt:Pt:Tiss:Qn

Note: All of the heavy metal measurements in hair, nails, and tissue should all be mass contents.

8157-0|Arsenic:MCnt:Pt:Nail:Qn

NCnt Number Content, number of component entities per unit mass of a sample (system).  
20771-2|Coliform bacteria:NCnt:Pt:Egg:Qn:Viability count

2. **Fraction (FR)**. Fraction of component A in a group of entities B, C, Y, N in system 1. By extension:

CFr Catalytic Fraction  
2536-1|Lactatedehydrogenase1/Lactatedehydrogenase.total:CFr:Pt:Ser:Plas:Qn:Electrophoresis  
9642-0|Creatine Kinase.BB/Creatine kinase.total:CFr:Pt:Ser:Plas:Qn

NFr Number Fraction  
10602-1|Spermatozoa.abnormal head/100 spermatozoa:NFr:Pt:Semen:Qn  
764-1|Neutrophils.band form/100 leukocytes:NFr:Pt:Bld:Qn:Manual count

MFr Mass Fraction  
2614-6|Methemoglobin/Hemoglobin.total:MFr:Pt:Bld:Qn

SFr Substance Fraction  
4546-8|Hemoglobin A/Hemoglobin.total:MFr:Pt:Bld:Qn

VFr Volume fraction.  
4545-0|Hematocrit:VFr:Pt:Bld:Qn:Spun

3. **Ratio (RTO)**. Ratio of component A to component B in system 1. By extension:

CCRto Catalytic Concentration Ratio  
2325-9|Gamma glutamyl transferase/Aspartate aminotransferase:CCRto:Pt:Ser:Plas:Qn

SCRto Substance Concentration Ratio  
2958-7|Sodium/Potassium:ScRto:Pt:Sweat:Qn

MCRto Mass Concentration Ratio  
2768-0|Phenylalanine/Tyrosine:MCrto:Pt:Ser:Plas:Qn

NRto Number Ratio  
11138-5|Myeloid cells/Erythroid cells:NRto:Pt:Bone mar:Qn

VelRto Velocity Ratio  
12022-0|Resistivity index:VelRto:Pt:Uterine artery.right:Qn:Doppler.calculated

VRatRto Volume Rate Ratio  
29462-9|Pulmonic flow/Systemic flow:VRatRto:Pt:Circulatory system.XXX:Qn:US.doppler

Ratio 1811-9|Amylase/Creatinine renal clearance:Ratio:24H:Urine:Qn

**Note:**

CSF/Serum Protein calculation is not a ratio, because the measured components are not in the same system. Its property type is relative mass concentration, RIMCnc (see below).

**Note:**

If the units of the denominator and numerator are both mass (e.g., mg/g), use MCcrto  
 13719-0|Carnitine/Creatinine:MCcrto:Pt:Urine:Qn

If the units of the denominator and numerator are both substance (e.g., mmol/mol) use ScRto  
 22695-1|Carnitine/Creatinine:ScRto:Pt:Urine:Qn

If the units of the denominator and numerator are different (mmol/g), use Ratio  
 17866-5|Carnitine/Creatinine:Ratio:Pt:Urine:Qn

**4. Relative (REL).** Relative amount of component A in system 1 compared to system 0. By extension:

REL should be used anywhere an actual measurement is divided by a measurement on a normal or control. It should also be used when a quotient is created by dividing a measured substance in Serum by the same substance measured in CSF, Urine, etc.

RelMCnc	Relative Mass Concentration (as noted previously) 2858-9 Protein.CSF/Protein.serum:RelMCnc:Pt:Ser+CSF:Qn 3235-9 Coagulation factor XII Ag actual/Normal:RelMCnc:Pt:PPP:Qn:Imm
RelTime	Relative time 3232-6 Coagulation factor XII activity actual/Normal:RelTime:Pt:PPP:Qn:Coag
RelCCnc	Relative Catalytic Concentration 28660-9 Plasminogen actual/Normal:RelCCnc:Pt:PPP:Qn:Chromo
RelRto	Relative Ratio 20450-3 Alpha-1-fetoprotein multiple of the median:RelRto:Pt:Ser/Plas:Qn
RelVol	Relative Volume 19853-1 Capacity.inspiratory.bs/Capacity.inspiratory.preop:RelVol:Pt:Respiratorysystem:Qn:Spirometry
RelVrat	Relative Volume Rate 20161-6 Voluntaryventilation.max^postbronchodilator/MVV:predicted:RelVrat:Pt:Respiratory system:Qn

**5. Cmplx.** Other divisions of one measurement by another that are not covered by the above rules should be classed as having Complex (Cmplx) properties, and the exact formula for deriving the quantity should be explicitly stated.

**6. ARBITRARY.** Arbitrary concentration of items. If we are not measuring the activity of an enzyme then the units of measure and properties are:

Possible Values	Property	Scale
Units/mL, IU/mL, etc.	ACnc	Qn
Units/gm, IU/gm, etc.	ACnt	Qn
Unit/min, IU/24hr, etc.	ARat	Qn
Unitless (Patient/Control)	AFr	Qn

When measuring presence/absence or ordering measures of a component, ACnc is also the correct property with scale of Ord

NOTE: If we are measuring the activity of an enzyme then the units of measure and properties are:

Possible Values	Property	Scale
IU/mL, Units/mL, etc.	CCnc	Qn
IU/gm, Units/gm, etc.	CCnt	Qn
IU/24hr, Unit/min, etc.	CRat	Qn
Unitless (Patient/Control)	CFr	Qn

**7. If the property is Titr then the scale is always Qn.**

For any X Ab or Ag:

Possible Values	Property	Scale
<1:2, 1:4, 1:8...	Titr	Qn

**8. For Any X Ab or Ag:**

Possible Values	Property	Scale
Neg, Indeterminate, Pos	ACnc	Ord
1+, 2+, 3+...	ACnc	Ord
<1:2, 1:4, 1:8...	Titr	Qn
Neg, 1:4, 1:8 ...	Titr	Qn
Neg, 0.90 ...	ACnc	Qn (EIA units)

9. For any intensive evaluation whose value comes from a finite set of unranked (independent) coded items the property will be Prid (or Type) and scale Nom. Prid is used in cases where the value set includes the option of reporting “none”, “not present”, etc. Type is used in cases where the result always specifies a value from the finite set. For extensive measures whose value comes from a finite set of unranked coded items, the property will be the extensive property, and the scale will be Nom.

Intensive Properties	Possible Values (coded)	Property	Scale
Organism Identified	E. coli, S. aureus, etc.	Prid	Nom
ABO Group	A, B, AB, O	Prid	Nom
Surgery (Dis. Summary)	Cholecystectomy, Appendectomy	Prid	Nom

Extensive Properties	Possible Values (coded)	Property	Scale
Urine Color	Amber, straw, etc.	Color	Nom
Urine Turbidity	Hazy, cloudy, opaque	Turbidity	Nom

10. For any intensive evaluation whose value comes from a finite set of unranked (independent) free text items (or a paragraph) the property will be Prid, or Find and scale Nar to indicate that the result is free text narrative. For extensive measures whose value comes from a finite set of unranked text items (or a paragraph), the property will be the extensive property, and the scale will be Nar.

Intensive Properties	Possible Values (text)	Property	Scale
Organism Identified	E. coli, S. aureus, etc.	Prid	Nar
ABO Group	A, B, AB, O	Prid	Nar
Surgery (Dis. Summary)	Cholecystectomy	Prid	Nar

Extensive Properties	Possible Values (text)	Property	Scale
Urine Color	Amber, straw, etc.	Color	Nar
Urine Turbidity	Hazy, cloudy, opaque	Turbidity	Nar

11. Imp is used to represent the property when the evaluation is a mental abstraction based on one a collection of measurements and or data. For example, if several measurements are made relative to immunoglobulin levels in Serum and CSF in a myasthenia gravis panel, and if by examining all of the evidence a pathologist decided that this pattern of findings represented active disease (which could be represented as a coded value), the result of the pathologist thought process would be represented as:

	<b>Possible Values (text)</b>	<b>Property</b>	<b>Scale</b>
Myasthenia Evaluation	No disease, chronic disease	Imp	Nom

If the pathologist evaluation is reported free text or a paragraph of information, the representation would be:

Myasthenia Evaluation	No disease, chronic disease	Imp	Nar
-----------------------	-----------------------------	-----	-----

12. Methods are only used to distinguish things that are identical in the other five LOINC fields but may differ because the sensitivity or specificity is different for the given methods.
13. Need to be careful in distinguishing end point detection method from property. For example, if sodium is measured using an ion specific electrode, the property is not a voltage difference. The voltage difference is just a method for indirectly measuring the sodium concentration. Concentration is the real property. Likewise, many antigens and antibodies are now measured using optical density as the detection method. However, the property we are really measuring is an arbitrary concentration (ACnc), not the optical density. If it is a ratio of optical densities (as with Gliadin Ab, Parvovirus B19 Ab, etc.) that are compared (patient value divided by a standard control), then the property should be ACRto (arbitrary concentration ratio).
14. ml/min/1.73sqM (Milliliters per min per 1.73 square meters BSA): Similar to the immediately preceding item. This result has the same property as if it had units of ml/min/sqM. The property of this measurement should be called "areic volume rate" (ArVRat).

## Appendix F - Acronyms used in LOINC

<b>Table 32: Acronyms used in LOINC</b>	
<b>Acronym</b>	<b>Meaning</b>
AC	Abdominal Circumference
ADL	Activities of Daily Living
AE	Anion Exchange protein
AP	Anterio-Posterior
APAD	AnteroPosterior Diameter of the Abdomen
AUT	Automated Ultrasound Testing
B2GPI	Beta 2 Glycoprotein 1
BD	Binocular Distance
BOR	Brachio-Oto-Renal
BPC	Biparietal Circumference
BPD	Biparietal diameter
CD	Cluster of differentiation
CDA	Congenital dyserythropoietic anaemia
CDB	Childhood Disability Benefits
cDNA	complementary DNA
CFst	Calorie Fast
CHAMPUS	Civilian Health and Medical Program of the Uniformed Services
Cine	Cinematographic
CNR1	Cannabinoid receptor 1
COC	Commission on Cancer
COPD	Chronic Obstructive Pulmonary Disease
CPT	Current Procedural Terminology
CRL	Crown-Rump Length
CSF	Cerebral spinal fluid
CW	Continuous wave
CyCD22	Cytoplasmic CD22
DBG	Donna Bennett-Goodspeed
DCIS	Ductal carcinoma in situ
DISIDA	Diisopropyliminodiacetic acid
DRG	Diagnostic Related Groups
DTPA	Diethylenetriamine pentaacetate
Dx	Diagnosis
EBV-LMP	Epstein Barr virus – latent membrane protein
ED	Emergency Department
EDD	Estimated Delivery Date
EEG	Electroencephalogra
EFW	Estimated Fetal Weight
EGD	Esophagogastro duodenoscopy
EKG	Electrocardiogram
EMS	Emergency Medical Service(s)
ENT	Ear, Nose Throat

ERCP	Endoscopic Retrograde Cholangiopancreatography
FL	Femur Length
FLACC	Face Legs Activity Cry Consolability
FNA	Fine needle
FTA	Fetal Trunk Area
GALOP	Gait disorder Autoantibody Late-age Onset Polyneuropathy
GSD	Gestational Sac Diameter
GSL	Gestational Sac Length
HC	Head Circumference
HCFA	Health Care Financing Administration
HIV	Human immunodeficiency virus
HIV-SSC	Sign and Symptom Check-List for Persons with HIV Disease
HL	Humerus Length
HLA	Human Leukocyte Antigen
HMPAO	Hexamethylpropyleneamine oxime
HTLV	Human T-cell Lymphotropic Virus
HWL	Height Width Length
ICD	International Classification of Diseases
ICD9	International Classification of Diseases, Ninth Revision
ICD9-CM	International Classification of Diseases, Ninth Revision, Clinical Modification
ICD-O	International Classification of Diseases for Oncology
ID	Intradermal
INR	International normalized ratio
IOD	Inter Ocular Distance
KUB	Kidney-Ureter-Bladder
LHON	Leber hereditary optic neuropathy
LOINC	Logical Observation Identifiers Names and Codes
LVOT	Left Ventricular Outflow Tract
LW	Landsteiner-Wiener
LWT	Length Width Thickness
MAA	Microalbumin aggregate albumin
MEMS	Medication Event Monitoring System
MERSTH	Medical Event Reporting System-Total Health System
MIB-1	Mindbomb homolog 1
MIBG	Metaiodobenzylguanidine
MIC	Minimum inhibitory concentration
MLC	Minimum lethal concentration
MLO	Mediolateral oblique
MMA	Macro aggregate albumin
MVV	Maximum Voluntary Ventilation
NAACCR	North American Association of Central Cancer Registries
Ng	Nasogastric
NPI	National Provider Identifier
OFD	Occipital-Frontal Diameter
O-I BPD	Outer to Inner Biparietal Diameter
OmpC	Outer membrane porin of E coli

O-O BPD	Outer to Outer Biparietal Diameter
O-O TD	Outer to Outer Tympanum Diameter
OOD	Outer Orbital Diameter
PA	Postero-Anterior
PCP	Primary Care Physician
PEG	Polyethylene Glycol
PHQ	Patient Health Questionnaire
PISA	Proximal Isovelocity Surface Area
PSR	Peridontal Screening and Recording
PYP	Pyrophosphate
QAM	Quality Audit Marker
QID	Four times a day
RAST	Radioallergosorbent test
RFC	Residual Functional Capacity
RFLP	Restriction fragment length polymorphism
RUG	Resource Utilization Groups
SAB	Streptoavidin-biotin
SBT	Sequence based typing
SC	Sulphur colloid
SCB	Sertoli cell barrier
SCL	Scleroderma
SEER	Surveillance Epidemiology and End Result
TAD	Transverse Abdominal Diameter
TC	Thoracic Circumference
TCD	Transverse Cerebellar Diameter
TD	Transaxial Diameter
TEC	Tubingen electric campimetry
TID	three times a day
TNM	Tumor, node, metastasis
TORCH	Toxoplasma, Rubella, Cytomegalovirus, Herpes Simplex Virus
TTD	Transverse Thoracic Diameter
TU	Tuberculin Units
VTI	Velocity Time Integral
VWF	von Willebrand Factor

## Appendix G - LOINC Committee Members

<b>Name</b>	<b>Organization</b>	<b>City, State/Province, Country</b>
Ray Aller	University Southern California Pathology	Vista, CA
John Baenziger	Indiana University Hospital	Indianapolis, IN
Suzanne Bakken	Columbia School of Nursing	New York, NY
Pam Banning	3M	West Linn, OR
Rita Barsoum	Kaiser Permanente	Pasadena, CA
James Barthel	H. Lee Moffitt Cancer Center	Tampa, FL
Dean Bidgood	Duke Medical Center	Durham, NC
Bruce Bray	University of Utah	Salt Lake City, UT
James Campbell	University of Nebraska	Omaha, NE
Jim Case	California Veterinary Diag Labs	Davis, CA
Jim Cimino	Columbia Presbyterian Med Center	New York, NY
Lori Carey	Canada Health Infoway	Saskatoon, SK, Canada
Robert Dolin	Mayo Foundation	Rochester, MN
James K Fleming	Laboratory Corp of America	Burlington, NC
Arden Forrey	University of Washington	Seattle, WA
Bill Francis	Augilent Technologies	Andover, MA
Pavla Frazier	University of Utah	Salt Lake City, UT
Alan Golichowski	Indiana Univ. Dept. of Medicine	Indianapolis, IN
Barry Gordon	C/NET Solutions	Berkeley, CA
Brian Griffin	Quest Diagnostics	Rutherford, NJ
Gil Hill	Hospital for Sick Children	Toronto, ON, Canada
Stan Huff	Intermountain Health Care	Salt Lake City, UT
Cindy Johns	LabCorp	Burlington, NC
William (Bill) Karitis	Department of Defense, U.S. Navy	Onley, MD
Ted Klein	Klein Consulting, Inc	Ridge, NY
Jeff Lamothe	USAF	Biloxi, MS
Lee Min Lau	3M HIS	Salt Lake City, UT
Diane Leland	Riley Hospital for Children	Indianapolis, IN
Pat Maloney	Quest Diagnostics	Teterboro, NJ
Doug Martin	Roudebush VA Medical Center	Indianapolis, IN
Susan Matney	Intermountain Health Care	Salt Lake City, UT
Ken McCaslin	Quest Diagnostics	Collegeville, PA
Clem McDonald	NLM Lister Hill National Center for Biomedical Communications	Bethesda, MD
Kathy Mercer	Regenstrief Institute	Indianapolis, IN
Deirdre O'Neill	National Medical Services Association	Willow Grove, PA
Judy Ozbolt	Vanderbilt University	Nashville, TN
Dan Pollock	Centers for Disease Control	Atlanta, GA
Rick Press	Oregon Health Sciences University	Portland, OR
Christine Raine	Parners Healthcare, Inc.	Brookline, MA
Angelo Rossi Mori	Instituto Tecnologie Biomediche	Rome, Italy
Jon Rosenblatt	Mayo Medical Laboratories	Rochester, MN
Shawn Shakib	3M HIS	Salt Lake City, UT
John Stelling	World Health Organization	Geneva, Switzerland
Steve Steindel	CDC	Atlanta, GA
Jeff Suico	Eli Lilly & Co.	Indianapolis, IN
Anders Thurin	University Hospital	Linkoping, Sweden
Wayne Tracy	Health Patterns, LLC	Overland Park, KS Daniel
Vreeman	Regenstrief Institute/IUSOM	Indianapolis, IN
Larry West	ARUP Laboratories	Salt Lake City, UT

Thomas White  
Warren Williams  
Pat Wilson

New York State Office of Mental Health  
CDC  
3M HIS

New York, NY  
Atlanta, GA  
Salt Lake City, UT

## LITERATURE CITED

- <sup>1</sup> Forrey AW, McDonald CJ, DeMoor G, Huff SM, Leavelle D, Leland D, Fiers T, Charles L, Stalling F, Tullis A, et. al. The logical observation identifier names and codes (LOINC) database: A public use set of codes and names for electronic reporting of clinical laboratory results. *Clinical Chemistry* 1996;42:81-90.
- <sup>2</sup> McDonald CJ, Huff SM, Suico JG, Hill G, Leavelle D, Aller R, Forrey A, Mercer K, DeMoor G, Hook J, Williams W, Case J, Maloney P. LOINC, a universal standard for identifying laboratory observations: A 5-Year update. *Clinical Chemistry* 2003;49:624-633.
- <sup>3</sup> ASTM E1238-94. Standard Specification for Transferring Clinical Observations Between Independent Computer Systems. Philadelphia: American Society for Testing Materials; 1994.
- <sup>4</sup> Health Level Seven. An application protocol for electronic data exchange in healthcare environments. Version 2.3. Ann Arbor, MI: Health Level Seven, Inc.; 1997.
- <sup>5</sup> McDonald CJ, Park BH, Blevins L. Grocers, physicians, and electronic data processing. *AMA Continuing Medical Education Newsletter* 1983;1:5-8.
- <sup>6</sup> International Union of Pure and Applied Chemistry/International Federation of Clinical Chemistry. The Silver Book: Compendium of terminology and nomenclature of properties in clinical laboratory sciences. Oxford: Blackwell Scientific Publishers; 1995.
- <sup>7</sup> Henry JB. *Clinical Diagnosis and Management by Laboratory Methods*. Philadelphia: W.B. Saunders; 1994.
- <sup>8</sup> Burtis CA, Ashwood ER (editors). *Tietz Textbook of Clinical Chemistry*, 2nd ed. Philadelphia: W.B. Saunders; 1994.
- <sup>9</sup> McDonald CJ, Tierney WM. Computer-stored medical records. *JAMA* 1988;259:3433-40.
- <sup>10</sup> Dick AS, Steen EB (editors). *The computer based patient record*. Washington DC: National Academy Press; 1991.
- <sup>11</sup> McDonald CJ, Overhage JM, Dexter P, Takesue BY, Dwyer DM. A Framework for capturing clinical data sets from computerized sources. *Ann Intern Med* 1997;8:675-682.
- <sup>12</sup> ASTM E1238-94. Standard Specification for Transferring Clinical Observations Between Independent Computer Systems. Philadelphia: American Society for Testing Materials; 1994.
- <sup>13</sup> *Ibid.*
- <sup>14</sup> Henry JB. *Clinical Diagnosis and Management by Laboratory Methods*. Philadelphia: W.B. Saunders; 1994.
- <sup>15</sup> International Union of Pure and Applied Chemistry/International Federation of Clinical Chemistry. The Silver Book: Compendium of terminology and nomenclature of properties in clinical laboratory sciences. Oxford: Blackwell Scientific Publishers; 1995.
- <sup>16</sup> Euclides Foundation International. EUCLIDES Laboratory Investigation Codes. Available from Dr. Georges DeMoor, Euclides Foundation International nv, Excelsioriaan 4A, B-1930, Zaventem, Belgium. Phone: 32 2 720 90 60.
- <sup>17</sup> Mahon CR, Manuselis G (editors). *Textbook of Diagnostic Microbiology*. Philadelphia: W.B. Saunders; 1995.
- <sup>18</sup> Walker RH. *American Association of Blood Banks Technical Manual*. 11th ed. Bethesda, MD: American Association of Blood Banks, 1993.
- <sup>19</sup> Euzéby JP. List of bacterial names with standing in nomenclature: a folder available on the internet. *Int J Syst Bacteriol* 1997;47:590-592. (List of prokaryotic names with standing in nomenclature. [Update 2008 May 2, cited 2008 June]. Available from: <http://www.bacterio.net>).
- <sup>20</sup> Index Virum, [Internet]. New York: New York, Catalogue of Viruses Version 2, based on the 6th ICTV Report; c2002 [Updated 2007 July; cited 2008 June]. Available from: <http://phene.cpmc.columbia.edu/>.
- <sup>21</sup> International Union of Pure and Applied Chemistry/International Federation of Clinical Chemistry. The Silver Book: Compendium of terminology and nomenclature of properties in clinical laboratory sciences. Oxford: Blackwell Scientific Publishers; 1995.
- <sup>22</sup> Stevens SS. Measurement, statistics, and the chemapiric view. Like the faces of Janus, science looks two ways – toward schematics and empirics. *Science* 1968;161:849-856.
- <sup>23</sup> Tang YW, Procop GW, Persing DH. Molecular diagnostics of infectious diseases. *Clin Chem* 1997;11:2021-2038.
- <sup>24</sup> Walker RH. *American Association of Blood Banks Technical Manual*. 11th ed. Bethesda, MD: American Association of Blood Banks, 1993.
- <sup>25</sup> Antonarakis, SE, and the Nomenclature Working Group. Recommendations for a nomenclature system for human gene mutations. *Human Mutation* 1998;11:1-3.
- <sup>26</sup> Beaudet AL, Tsui LC. A suggested nomenclature for designating mutations. *Human Mutation* 1993;2:245-248.
- <sup>27</sup> ImmunoCAP Allergens [Internet]. Uppsala(Sweden): ImmunoCAP; c2006 [updated 2007 November 2; cited 2008 June]. Available from: <http://www.immunocapinvitrosight.com/>.
- <sup>28</sup> Ballinger PW, Frank ED. 10th ed. Vol. 3 Merrill's atlas of radiographic positions and radiographic procedures. St. Louis: Elsevier Mosby; 2003.
- <sup>29</sup> Dorland's Illustrated Medical Dictionary. 30th ed. Philadelphia: Saunders/Elsevier; 2003.
- <sup>30</sup> Bakken S, et al. Evaluation of Clinical LOINC (Logical Identifiers, Names, and Codes) for Terminology Model for Standardized Assessment of Measures. *JAMIA* 7:2000; 529-538.